



**HAL**  
open science

## Signed tropical halfspaces and convexity

Georg Loho, Mateusz Skomra

► **To cite this version:**

| Georg Loho, Mateusz Skomra. Signed tropical halfspaces and convexity. 2022. hal-03765509

**HAL Id: hal-03765509**

**<https://laas.hal.science/hal-03765509>**

Preprint submitted on 31 Aug 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# SIGNED TROPICAL HALFSPACES AND CONVEXITY

GEORG LOHO AND MATEUSZ SKOMRA

ABSTRACT. We extend the fundamentals for tropical convexity beyond the tropically positive orthant expanding the theory developed by Loho and Végé (ITCS 2020). We study two notions of convexity for signed tropical numbers called *TO-convexity* (formerly ‘signed tropical convexity’) and the novel notion *TC-convexity*.

We derive several separation results for TO-convexity and TC-convexity. A key ingredient is a thorough understanding of TC-hemispace – those TC-convex sets whose complement is also TC-convex. Furthermore, we use new insights in the interplay between convexity over Puiseux series and its signed valuation.

Remarkably, TC-convexity can be seen as a natural convexity notion for representing oriented matroids as it arises from a generalization of the composition operation of vectors in an oriented matroid.

## 1. INTRODUCTION

Convexity is a powerful structure which is often behind the existence of efficient algorithms. In this spirit, tropical convexity, arising from classical convexity by replacing addition with maximization and multiplication by addition, found several applications from optimization [11], phylogenetics [54] and machine learning [43]. While tropical convexity was mainly considered with a non-negativity constraint for a long time, the recent paper [42] introduced a notion of signed tropical convexity to overcome this restriction. In this paper, we extend the fundamentals of signed tropical convexity and exhibit a slightly different version of signed tropical convexity.

The crucial building blocks of the two versions of signed tropical convexity are open and closed tropical halfspaces. We coin the name *TO-convexity* (tropical open convexity) for the convexity in which the hull of finitely many points equals the intersection of the *open* tropical halfspaces containing them. Additionally, we define *TC-convexity* (tropical closed convexity) in such a way that the hull of finitely many points equals the intersection of the *closed* tropical halfspaces containing them. For technical reasons, we actually introduce it via the hull of only two points first (Definition 3.5) and give the characterization for bigger sets by closed tropical halfspaces as one of our main results.

Our main result is Theorem 6.1 which shows that the definition via hulls of only two points is actually equivalent to the definition via intersection of closed tropical halfspaces. This gives an analog of the hyperplane separation theorem for TC-convexity. To arrive at this statement, we need to gather extensive insights into the structure of TC-convex sets and their lifts to Puiseux series. We make explicit use of the connection between separation of sets over Puiseux series and their signed valuation established in Theorem 4.10. We use this to strengthen the separation

results which we derive directly for TC-convex sets. The main insights for those is the fundamental elimination property of TC-convexity given in Theorem 5.2. To combine this with the separation properties for lifts, we investigate TC-hemispace – those TC-convex sets whose complement is also TC-convex. We show that they are nearly tropical halfspaces in Theorem 5.10 and provide several structural insights on their boundary. Along the way, we also give new separation results for TO-convex sets including the Pasch & Kakutani properties (Theorem 3.3). Furthermore, we derive Carathéodory-type results for TC-convexity. We finish with Minkowski–Weyl theorems for finitely generated TC-convex sets (Theorem 6.3 and Theorem 6.4).

**Comparison of TO-convexity and TC-convexity.** From the viewpoint of abstract convexity [51], TO-convexity is generated by open tropical halfspaces and TC-convexity is generated by closed tropical halfspaces. This yields that TO-convex sets are also TC-convex. While TO-convex sets still behave rather well with the algebraic operations, as TO-convex hull can be written using (tropical) convex combinations with hyperoperations, this structure is lost for TC-convexity.

To describe the latter, we use a non-commutative addition, which we call *left sum* (Definition 2.11). It generalizes the composition of vectors in an oriented matroid. In this way, this operation already appeared in work on Bergman fans for matroid over hyperfields generalizing composition of sign vectors of an oriented matroid [23, 15]. We will elaborate more about the implication of this connection in upcoming work.

Another remarkable difference is the behavior with respect to lifts to Puiseux series. In [42, Theorem 3.14], it was shown that the TO-convex hull of finitely many points arises as union of the signed valuations of the convex hulls ranging over all lifts. We show in Corollary 6.2, that the TC-convex hull arises as the intersection of these lifts.

**Motivation.** Our work is motivated by the need for a better structural basis underlying (recent) applications of tropical convexity. New insights on the complexity of classical linear programming based on tropical geometry have been a great success story in recent years [5, 10, 11, 14]. The point of origin for these advances is the tropicalization of linear programs – where one always had to impose additional nonnegativity constraints. A thorough foundation of signed tropical convexity will allow to study the signed tropicalization of general linear programs.

Furthermore, tropical convexity has an intimate connection with mean payoff games as the feasibility problem for tropical inequality systems (restricted to the tropical nonnegative orthant) is equivalent to mean payoff games [2]. The latter have an intriguing complexity status in the intersection of NP and co-NP while no polynomial-time algorithm is known [57]; see the recent paper [26] for a good overview including the flourishing advances on the subclass of parity games. Studying the polar of a tropical polyhedron naturally leads to tropically convex sets without a nonnegativity constraint which was modeled by pairs of nonnegative numbers in former work [12]. We think that signed tropical convexity may enrich the insights in the structure of these games.

Yet another motivation comes from interest in signed tropicalization of semialgebraic sets [36]. The study of unsigned tropicalizations of semialgebraic sets already lead to a fruitful connection between stochastic games and (non-Archimedean) semi-definite programs [7]. In particular, the analysis of tropical cones arising in this

work resulted in a recent universal complexity bounds on value iteration for a large class of games in [6]. Our framework allows to capture also other classes of optimization problems where the nonnegativity condition is a priori not satisfied.

Finally, already in [42], it was demonstrated that signed tropical linear inequality systems with unit coefficients are the same as Boolean formulas. In this sense, more general signed tropical linear inequality systems form a ‘quantitative’ generalization of Boolean formulas. Therefore, the feasibility problem for linear systems over signed tropical numbers is a natural generalization of SAT. Note that the solution sets of these systems are actually not TO-convex. However, they are exactly the finitely generated TC-convex sets.

**Related work.** We refer to the book [37] for a general overview on (unsigned) tropical convexity. Different versions of separation theorems in the tropical convexity have been obtained by numerous authors [56, 48, 25, 28, 22, 33]. Likewise, the Carathéodory theorem for tropical convexity has been discovered independently in different works [34, 21, 28]. The tropical analogue of the Minkowski–Weyl theorem for was proven in [31]. In order to prove our main separation theorem, we rely on works on abstract convexity, which were already applied to the usual tropical convexity in [35]. Furthermore, our proof requires to give a partial characterization of signed tropical hemispaces. A full characterization of tropical hemispaces in one orthant is given in [22, 39, 30].

The link between the tropical convexity and convexity over Puiseux series was established in the work [29] which studies the tropicalization of polyhedra. This was later generalized to spectrahedra in [55, 13] and to convex semialgebraic sets in [8]. The tropicalizations of general semialgebraic sets are studied in [4] and this study is extended to signed tropicalizations in [36] and [13, Section 4].

Signed tropical numbers first appeared in the context of the symmetrized semiring [1]. The idea that tropical convexity could be extended to signed tropical numbers appeared already in [21, 20], where two such extensions (different than the ones considered here) are introduced. Our work is most closely related to the paper [42] which introduces and studies the TO-convexity.

## 2. PRELIMINARIES ON SIGNED TROPICAL NUMBERS

We give a brief overview of necessary notions related to signed numbers; for more see [3, 1, 42]. The signed tropical numbers  $\mathbb{T}_\pm$  are obtained by glueing two copies of  $(\mathbb{R} \cup \{-\infty\})$  at the tropical zero element  $\mathbb{O} = -\infty$  giving rise to the *non-negative tropical numbers*  $\mathbb{T}_{\geq \mathbb{O}} = \mathbb{R} \cup \{\mathbb{O}\}$  and the *non-positive tropical numbers*  $\mathbb{T}_{\leq \mathbb{O}} = \{\ominus x \mid x \in \mathbb{R}\} \cup \{\mathbb{O}\}$ .

The signed numbers can be extended to the *symmetrized semiring*  $\mathbb{S}$ , which forms a semiring containing  $\mathbb{T}_\pm$ , by attaching *balanced numbers*  $\mathbb{T}_\bullet = \{\bullet x \mid x \in \mathbb{R}\} \cup \{\mathbb{O}\}$ . We will often use the norm  $|\cdot|$  on  $\mathbb{T}_\pm$  which maps each element of  $\mathbb{T}_{\geq \mathbb{O}}$  to itself, removes the  $\ominus$  sign of an element in  $\mathbb{T}_{\leq \mathbb{O}}$  and the  $\bullet$  from an element in  $\mathbb{T}_\bullet$ . This is complemented by the map  $\text{tsgn}$  from  $\mathbb{S}$  to  $\{\oplus, \ominus, \bullet, \mathbb{O}\}$  keeping only the sign information.

Throughout, we use the notation  $[d] = \{1, 2, \dots, d\}$  and  $[d]_0 = [d] \cup \{0\}$ . For a vector  $z \in \mathbb{T}_\pm^d$  we denote its support, positive support and negative support,

respectively, by

$$\begin{aligned} \text{supp}(z) &= \{i \in [d] \mid z_i \neq \mathbf{0}\} \\ \text{supp}^\oplus(z) &= \{i \in [d] \mid z_i \in \mathbb{T}_{>\mathbf{0}}\} \\ \text{supp}^\ominus(z) &= \{i \in [d] \mid z_i \neq \mathbb{T}_{<\mathbf{0}}\} . \end{aligned} \quad (1)$$

For  $x, y \in \mathbb{S}$ , the addition is defined by

$$x \oplus y = \begin{cases} \text{argmax}_{x,y}(|x|, |y|) & \text{if } |\chi| = 1 \\ \bullet \text{argmax}_{x,y}(|x|, |y|) & \text{else} . \end{cases} \quad (2)$$

where  $\chi = \{\text{tsgn}(\xi) \mid \xi \in (\text{argmax}(|x|, |y|))\}$ . The multiplication is given by

$$x \odot y = (\text{tsgn}(x) * \text{tsgn}(y)) (|x| + |y|) , \quad (3)$$

where the  $*$ -multiplication table is the usual multiplication of  $\{-1, 1, 0\}$  for  $\{\ominus, \oplus, \bullet\}$  with the additional specialty that multiplication with  $\mathbf{0}$  yields  $\mathbf{0}$ . The operations  $\oplus$  and  $\odot$  extend to vectors and matrices componentwise.

We use  $\oplus, \ominus$  and  $\bullet$  also as unary operations on  $\mathbb{S}$  with  $\ominus\ominus = \oplus, \bullet\ominus = \bullet\bullet = \ominus\bullet = \bullet$  and where  $\oplus$  just acts as identity. The fact that  $\mathbb{S}$  with these operations forms a commutative semiring justifies the name introduced above. We also point out that balanced numbers in the symmetrized semiring are equivalent to the use of a multivalued addition as in the theory of hyperfields [27, 17]. More precisely, the addition in the symmetrized semiring is equivalent to the multivalued addition in the *real tropical hyperfield* discussed in [36, 53].

*Example 2.1.* One has  $2 \odot \ominus 1 = \ominus 3$ ,  $(0 \oplus \ominus 0) \odot \ominus - 1 = \bullet 0 \odot \ominus - 1 = \bullet - 1$ ,  $-1 \odot -1 = -2$ ,  $\ominus 1 \odot \ominus 1 = 2$ .

We recall some relations which serve to order the symmetrized semiring  $\mathbb{S}$ . For  $x, y \in \mathbb{S}$  we set

$$\begin{aligned} x > y &\Leftrightarrow x \ominus y \in \mathbb{T}_{>\mathbf{0}} \\ x \vDash y &\Leftrightarrow x \ominus y \in \mathbb{T}_{>\mathbf{0}} \cup \mathbb{T}_\bullet \\ x \geq y &\Leftrightarrow x > y \vee x = y \end{aligned} \quad (4)$$

For  $a \in \mathbb{S}$ , we set

$$\mathcal{U}(a) = \begin{cases} [\ominus|a|, |a|] = \{x \in \mathbb{T}_\pm \mid \ominus|a| \leq x \leq |a|\} & \text{for } a \in \mathbb{T}_\bullet \\ \{a\} & \text{else} \end{cases} . \quad (5)$$

We extend this to vectors by setting  $\mathcal{U}(v) = \prod_{i \in [d]} \mathcal{U}(v_i)$ .

*Example 2.2.* One has  $2 > \ominus 3$  but  $\bullet 4$  and  $\ominus 3$  are incomparable via ' $>$ ' since  $\bullet 4 \ominus \ominus 3 = \bullet 4 \oplus 3 = \bullet 4 = \ominus 4 \bullet 4$ . Though it holds  $\bullet 4 \vDash \ominus 3$  and  $\ominus 3 \vDash \bullet 4$  and  $3 \vDash \bullet 4$  showing, e.g., that ' $\vDash$ ' is not anti-symmetric.

The signed tropical numbers are equipped with the order topology induced by the strict order  $<$ . With this,  $\mathbb{T}_\pm$  is homeomorphic to  $\mathbb{R}$  with the usual order topology via

$$\text{slog}: \mathbb{R} \rightarrow \mathbb{T}_\pm, \quad \text{slog}(x) = \begin{cases} \log(x) & \text{for } x > 0 \\ \ominus \log(|x|) & \text{for } x < 0 \\ \mathbf{0} & \text{for } x = 0 \end{cases} .$$

This extends to  $\mathbb{T}_{\pm}^d$  via the product topology, in particular  $\mathbb{T}_{\pm}^d$  is homeomorphic to  $\mathbb{R}^d$ . Hence, we can use all topological properties of  $\mathbb{R}^d$  also for vectors of signed tropical numbers.

Recall that a partial order on some set  $S$  is *dense* if for every two elements  $x, y \in S$  such that  $x < y$  there exists  $z \in S$  that satisfies  $x < z < y$ . With this, the order  $<$  on  $\mathbb{T}_{\pm}$  is dense.

**2.1. Halfspaces.** The different versions of halfspaces form the building blocks for our convexity notions.




FIGURE 1. An open and a closed signed tropical halfspace.

**Definition 2.3.** For a vector  $(a_0, a_1, \dots, a_d) \in \mathbb{T}_{\pm}^{d+1}$  such that  $(a_1, \dots, a_d) \neq \mathbf{0}$  we define the *signed (affine) tropical hyperplane* by

$$\mathcal{H}(a) = \left\{ x \in \mathbb{T}_{\pm}^d \mid a \odot \begin{pmatrix} 0 \\ x \end{pmatrix} \in \mathbb{T}_{\bullet} \right\}, \quad (6)$$

the *open signed (affine) tropical halfspace* by

$$\mathcal{H}^+(a) = \left\{ x \in \mathbb{T}_{\pm}^d \mid a \odot \begin{pmatrix} 0 \\ x \end{pmatrix} \in \mathbb{T}_{>\mathbf{0}} \right\}, \quad (7)$$

the *closed signed (affine) tropical halfspace* by

$$\overline{\mathcal{H}}^+(a) = \left\{ x \in \mathbb{T}_{\pm}^d \mid a \odot \begin{pmatrix} 0 \\ x \end{pmatrix} \in \mathbb{T}_{>\mathbf{0}} \cup \mathbb{T}_{\bullet} \right\}, \quad (8)$$

and the *semi-closed signed (affine) tropical halfspace* by

$$\tilde{\mathcal{H}}^+(a) = \left\{ x \in \mathbb{T}_{\pm}^d \mid a \odot \begin{pmatrix} 0 \\ x \end{pmatrix} \in \mathbb{T}_{\geq\mathbf{0}} \right\}. \quad (9)$$

If  $a_0 = \mathbf{0}$ , we call a tropical halfspaces *linear* instead of affine.

We denote  $\mathcal{H}^-(a) = \mathcal{H}^+(\ominus a)$  and  $\overline{\mathcal{H}}^-(a) = \overline{\mathcal{H}}^+(\ominus a)$ . Furthermore, given  $J \subseteq [d]$ , we say that a tropical halfspace  $\overline{\mathcal{H}}^+(a)$  is of *type*  $J$  if  $J = \{i \geq 1 \mid a_i \in \mathbb{T}_{>\mathbf{0}}\}$ .

To express  $\overline{\mathcal{H}}^+(a)$  using a more classical notation, we recall that  $\max \emptyset = -\infty$ . Then, we have  $x \in \overline{\mathcal{H}}^+(a)$  if and only if

$$\max(a_0^+, \max_{\text{tsgn}(x_i)=\text{tsgn}(a_i)} (|a_i| + |x_i|)) \geq \max(a_0^-, \max_{\text{tsgn}(x_i) \neq \text{tsgn}(a_i)} (|a_i| + |x_i|)), \quad (10)$$

where  $(a_0^+, a_0^-) = (a_0, -\infty)$  for  $a_0 \in \mathbb{T}_{\geq\mathbf{0}}$  and  $(a_0^+, a_0^-) = (-\infty, |a_0|)$  otherwise. We have  $x \in \mathcal{H}^+(a)$  if and only if the inequality in (10) is strict. The following lemma gathers the basic topological properties of tropical halfspaces.

**Lemma 2.4.** *The set  $\overline{\mathcal{H}^+}(a)$  is closed, its interior is equal to  $\mathcal{H}^+(a)$ , and the closure of  $\mathcal{H}^+(a)$  is equal to  $\overline{\mathcal{H}^+}(a)$ .*

*Proof.* Equation (10) shows that the restriction of  $\overline{\mathcal{H}^+}(a)$  to any closed orthant of  $\mathbb{T}_\pm^d$  is closed. Therefore,  $\overline{\mathcal{H}^+}(a)$  is closed and  $\mathcal{H}^+(a) = \mathbb{T}_\pm^d \setminus \overline{\mathcal{H}^-}(a)$  is open. Also, [42, Lemma 5.2] shows that  $\overline{\mathcal{H}^+}(a)$  is the closure of  $\mathcal{H}^+(a)$ . In particular,  $\mathcal{H}^+(a)$  is included in the interior of  $\overline{\mathcal{H}^+}(a)$ . To finish, we note that the interior of  $\overline{\mathcal{H}^+}(a)$  cannot contain any point that belongs to  $\mathcal{H}^-(a) \subseteq \overline{\mathcal{H}^-}(a)$  because  $\overline{\mathcal{H}^-}(a)$  is the closure of  $\mathcal{H}^-(a)$  and  $\overline{\mathcal{H}^+}(a) \cap \mathcal{H}^-(a) = \emptyset$ .  $\square$

**2.2. Basic calculations in  $\mathbb{S}$ .** We collect and extend basic properties from [42].

**Lemma 2.5** ([42, Lemma 2.6]). *Let  $a, b, c, d \in \mathbb{S}$ .*

- (a)  $a \oplus c \vDash b \Leftrightarrow a \vDash b \ominus c$
- (b)  $a \vDash b \wedge c \vDash d \Rightarrow a \oplus c \vDash b \oplus d$ .
- (c) *If  $c \in \mathbb{T}_\pm$ , then  $b \vDash c$  and  $c \vDash a$  imply  $b \vDash a$ .*
- (d)  $a \vDash b$  implies  $c \odot a \oplus d \vDash c \odot b \oplus d$  for  $c \in \mathbb{T}_{\geq 0}$  and  $c \odot b \oplus d \vDash c \odot a \oplus d$  for  $c \in \mathbb{T}_{\leq 0}$ .

**Lemma 2.6.** *Let  $a, b, c \in \mathbb{S}$ . Then,*

$$a \oplus c \vDash \mathbf{0} \wedge b \oplus c \vDash \mathbf{0} \Rightarrow a \oplus b \oplus c \vDash \mathbf{0} \wedge a \oplus b \ominus c \vDash \mathbf{0}.$$

*Proof.* We distinguish two cases.

**Case 1** ( $c \in \mathbb{T}_\bullet$ ) Adding the equivalent relations  $a \vDash c$  and  $b \vDash \ominus c$  side-wise yields  $a \oplus b \vDash \bullet c$ . This is equivalent to  $a \oplus b \bullet c \vDash \mathbf{0}$  and shows the required as in this case  $\bullet c = \ominus c = c$ .

**Case 2** ( $c \in \mathbb{T}_\pm$ ) Without loss of generality, we can assume that  $c \geq \mathbf{0}$ . With  $a \vDash c$  this implies  $|a| \geq |c|$ . Combining  $a \vDash c$  and  $c \vDash \ominus b$  yields  $a \oplus b \vDash \mathbf{0}$  in this case. With

$$|a \oplus b| = |a| \oplus |b| \geq |a| \geq |c|$$

we get  $a \oplus b \vDash c \geq \ominus c$  by checking the two possibilities  $a \oplus b \in \mathbb{T}_\pm$  or  $a \oplus b \in \mathbb{T}_\bullet$ .  $\square$

**Lemma 2.7.** *Let  $u, v, w \in \mathbb{T}_\pm$ ,  $q \in \mathcal{U}(v \oplus w)$ ,  $p \in \mathcal{U}(u \oplus v)$ .*

- (a)  $\mathcal{U}(u \oplus q) \subseteq \mathcal{U}(u \oplus v \oplus w)$  and  $\mathcal{U}(p \oplus w) \subseteq \mathcal{U}(u \oplus v \oplus w)$ ,
- (b)  $\mathcal{U}(u \oplus q) \cap \mathcal{U}(p \oplus w) \neq \emptyset$ .

*Proof.* For (a): By commutativity, it suffices to prove the first inclusion. For  $v \oplus w \in \mathbb{T}_\pm$ , we have  $q = v \oplus w$  and therefore  $\mathcal{U}(u \oplus q) = \mathcal{U}(u \oplus v \oplus w)$ . Otherwise, the claim follows from the definition of  $\mathcal{U}(\cdot)$  using

$$|u \oplus q| = |u| \oplus |q| \leq |u| \oplus |u \oplus w| = |u \oplus v \oplus w|$$

by distinguishing  $|u| > |v \oplus w|$  and  $|u| \leq |v \oplus w|$ .

For (b):

**Case 1** ( $\mathcal{U}(v \oplus w)$  or  $\mathcal{U}(u \oplus v)$  is singleton.) Assume without loss of generality that  $\mathcal{U}(v \oplus w)$  is a singleton. This implies  $\mathcal{U}(u \oplus q) = \mathcal{U}(u \oplus v \oplus w) \supseteq \mathcal{U}(p \oplus w)$ , where the inclusion follows from (a).

**Case 2** ( $v \oplus w$  and  $u \oplus v$  are balanced.) Here, we have  $u = \ominus v = w$ . This means we have either  $p = \ominus w$  and, hence,  $\mathcal{U}(p \oplus w) = \mathcal{U}(u \oplus v \oplus w)$ , or  $p \oplus w = w = u$ . The same applies to  $q$  and  $u$ . Therefore, in each of the four combinations of the two possibilities, one has a non-empty intersection.  $\square$

As  $\mathcal{U}(\cdot)$  is defined componentwise, we obtain a higher-dimensional extension of Lemma 2.7(b).

**Corollary 2.8.** *For  $u, v, w \in \mathbb{T}_\pm^d$  and  $q \in \mathcal{U}(v \oplus w)$ ,  $p \in \mathcal{U}(u \oplus v)$ , one gets  $\mathcal{U}(u \oplus q) \cap \mathcal{U}(p \oplus w) \neq \emptyset$ .*

**Lemma 2.9** ([42, Lemma 3.5(b)]). *If  $a \in \mathcal{U}(x)$ ,  $b \in \mathcal{U}(y)$ , and  $c \in \mathbb{T}_\pm$ , then  $\mathcal{U}(c \odot a \oplus b) \subseteq \mathcal{U}(c \odot x \oplus y)$ .*

*Observation 2.10.* Let  $a, b, \mu \in \mathbb{T}_\pm$  and  $s, t \in \mathbb{S}$  with  $a \in \mathcal{U}(s)$  and  $b \in \mathcal{U}(t)$ . Then  $\mu \odot a \in \mathcal{U}(\mu \odot s)$  and  $a \oplus b \in \mathcal{U}(s \oplus t)$ .

**2.3. Left sum.** The following notion of one-sided addition of tropical signed numbers is crucial for our study of convexity over signed tropical numbers.

**Definition 2.11.** Given two tropical numbers  $x, y \in \mathbb{S}$  we define their *left sum* as

$$x \triangleleft y = \begin{cases} x & \text{if } |x| = |y|, \\ x \oplus y & \text{otherwise.} \end{cases}$$

We extend this definition to vectors  $x, y \in \mathbb{S}^d$  by putting  $(x \triangleleft y)_k = x_k \triangleleft y_k$  for every  $k \in [d]$ .

*Remark 2.12.* For the special case of vectors with entries in  $\{\mathbb{O}, 0, \ominus 0\}$ , the left sum operation is just the *composition* of sign vectors in an oriented matroid [19]. The generalization to signed vectors with real entries already appears for the description of *real Bergman fans* [23] and even more generally in the context of matroids over tracts [15, Section 6.2]. From this point of view, later on in Proposition 3.17 we will see an extension of the elimination property of oriented matroids. We will discuss this connection further in upcoming work.

The left sum operation is not commutative as one can see on the example  $0 \oplus (\ominus 0) = 0 \neq \ominus 0 = (\ominus 0) \oplus 0$ . Nevertheless, it is associative and compatible with the order on  $\mathbb{T}_\pm$ .

*Observation 2.13.* For every  $x, y, z \in \mathbb{T}_\pm^d$  we have

- (a)  $(x \triangleleft y) \triangleleft z = x \triangleleft (y \triangleleft z)$
- (b)  $|x \triangleleft y| = |x| \oplus |y|$ .

**Lemma 2.14.** *Let  $a_1, b_1, a_2, b_2 \in \mathbb{T}_\pm$ .*

- (a) *If  $a_1 \leq b_1$  and  $a_2 \leq b_2$ , then  $a_1 \triangleleft a_2 \leq b_1 \triangleleft b_2$ .*
- (b) *if  $a_1 < b_1$  and  $a_2 < b_2$ , then  $a_1 \triangleleft a_2 < b_1 \triangleleft b_2$ .*
- (c) *if  $a_1 \geq b_1$  and  $a_2 \geq b_2$ , then  $a_1 \triangleleft a_2 \geq b_1 \triangleleft b_2$*
- (d) *if  $a_1 > b_1$  and  $a_2 > b_2$ , then  $a_1 \triangleleft a_2 > b_1 \triangleleft b_2$ .*

*Proof.* It is enough to prove (a) and (b) as (c),(d) follow by replacing  $(a_1, a_2, b_1, b_2)$  with  $(\ominus a_1, \ominus a_2, \ominus b_1, \ominus b_2)$ . Hence, we suppose that  $a_1 \leq b_1$  and  $a_2 \leq b_2$  (or  $a_1 < b_1$  and  $a_2 < b_2$ ). Let  $x = a_1 \triangleleft a_2$  and  $y = b_1 \triangleleft b_2$ .

If  $(x, y) \in \{(a_1, b_1), (a_2, b_2)\}$ , then the claim is trivial.

If  $(x, y) = (a_1, b_2)$ , then we have  $|b_2| \geq |b_1|$  and  $|a_1| \geq |a_2|$ . If  $b_2 \geq \mathbb{O}$ , then  $b_2 \geq b_1 \geq a_2$  (or  $b_2 > a_1$  if  $a_1 < b_1$ ). If  $b_2 < \mathbb{O}$ , then  $a_2 \leq b_2 < \mathbb{O}$  implies that  $|a_1| \geq |a_2| \geq |b_2| \geq |b_1|$ . Therefore,  $a_1 \leq b_1$  implies that  $a_1 \leq \mathbb{O}$ . Since  $|a_1| \geq |a_2|$ , we get  $a_1 \leq a_2 \leq b_2$  (or  $a_1 < b_2$  if  $a_2 < b_2$ ).

If  $(x, y) = (a_2, b_1)$ , then we have  $|a_2| \geq |a_1|$ ,  $|b_1| \geq |b_2|$ , and an analogous proof as above applies.  $\square$



**Lemma 2.15.** Fix  $a, b \in \mathbb{T}_\pm$  with  $\text{tsgn}(a) = \text{tsgn}(b)$  and let  $x_1^{(1)}, \dots, x_n^{(1)} \in \mathbb{T}_\pm$ ,  $x_1^{(2)}, \dots, x_n^{(2)} \in \mathbb{T}_\pm$  be such that

$$a \leq x_1^{(1)} \triangleleft \dots \triangleleft x_n^{(1)} \quad \text{and} \quad b \leq x_1^{(2)} \triangleleft \dots \triangleleft x_n^{(2)}.$$

Then

$$a \oplus b \leq w_1 \triangleleft \dots \triangleleft w_n$$

for all  $w_1 \in \mathcal{U}(x_1^{(1)} \oplus x_1^{(2)}), \dots, w_n \in \mathcal{U}(x_n^{(1)} \oplus x_n^{(2)})$ .

By multiplying with  $\ominus 0$ , the implication also holds if one replaces ' $\leq$ ' by ' $\geq$ '.

*Proof.* Let  $p, q \in [n]$  be the smallest indices such that  $x_1^{(1)} \triangleleft \dots \triangleleft x_p^{(1)} = x_p^{(1)}$  and  $x_1^{(2)} \triangleleft \dots \triangleleft x_q^{(2)} = x_q^{(2)}$ . Then Lemma 2.14 implies that  $a \triangleleft b \leq x_p^{(1)} \triangleleft x_q^{(2)}$  and  $b \triangleleft a \leq x_q^{(2)} \triangleleft x_p^{(1)}$ . Due to  $\text{tsgn}(a) = \text{tsgn}(b)$ , this yields  $a \oplus b \leq \min(x_p^{(1)} \triangleleft x_q^{(2)}, x_q^{(2)} \triangleleft x_p^{(1)})$ .

By definition, for any  $s, t \in \mathbb{T}_\pm$ , the interval  $\mathcal{U}(s \oplus t)$  has the boundary points  $s \triangleleft t$  and  $t \triangleleft s$ . There is a choice of orders  $\tau_1, \dots, \tau_n \in \text{Sym}(2)$  which minimizes the expression  $\triangleleft (x_i^{(\tau_i(1))} \triangleleft x_i^{(\tau_i(2))})$ . By the former observation and Lemma 2.14 it is smaller than  $w_1 \triangleleft \dots \triangleleft w_n$  for all  $w_1 \in \mathcal{U}(x_1^{(1)} \oplus x_1^{(2)}), \dots, w_n \in \mathcal{U}(x_n^{(1)} \oplus x_n^{(2)})$ . Let  $x_j^{(\ell)}$  for  $j \in [n]$  and  $\ell \in \{1, 2\}$  be the entry which defines the value of this expression, which means the first summand with maximal absolute value. If  $\ell = 1$  then  $j = p$ , otherwise  $j = q$ . In either case, we get  $\triangleleft (x_i^{(\tau_i(1))} \triangleleft x_i^{(\tau_i(2))}) = x_p^{(1)} \triangleleft x_q^{(2)}$  or  $\triangleleft (x_i^{(\tau_i(1))} \triangleleft x_i^{(\tau_i(2))}) = x_q^{(2)} \triangleleft x_p^{(1)}$ . In particular, we obtain

$$w_1 \triangleleft \dots \triangleleft w_n \geq \triangleleft (x_i^{(\tau_i(1))} \triangleleft x_i^{(\tau_i(2))}) \geq \min(x_p^{(1)} \triangleleft x_q^{(2)}, x_q^{(2)} \triangleleft x_p^{(1)}) \geq a \oplus b. \quad \square$$

*Example 2.16.* Let  $a = 0$ ,  $b = \ominus 0$ ,  $x_1^{(1)} = \mathbf{0}$ ,  $x_2^{(1)} = 0$ ,  $x_1^{(2)} = \ominus 0$ ,  $x_2^{(2)} = \mathbf{0}$ ,  $w_1 = \ominus 0$ ,  $w_2 = 0$ . Then  $a \leq x_1^{(1)} \triangleleft x_2^{(1)}$ ,  $b \leq x_1^{(2)} \triangleleft x_2^{(2)}$ ,  $w_1 = x_1^{(1)} \oplus x_1^{(2)}$ ,  $w_2 = x_2^{(1)} \oplus x_2^{(2)}$ , but  $0 = a \triangleleft b > w_1 \triangleleft w_2 = \ominus 0$ . This example shows that the assumption  $\text{tsgn}(a) = \text{tsgn}(b)$  cannot be omitted in the previous lemma.

### 3. FLAVORS OF SIGNED CONVEXITY

In [42], a notion of convexity for signed tropical numbers was introduced. We revisit this notion under the name *TO-convexity* and give several new insights. Furthermore, we establish a second notion of convexity for signed tropical numbers in an equally natural way.

Recall that a *cone* is a set  $X \subseteq \mathbb{T}_\pm^d$  such that  $\lambda \odot X \subseteq X$  for all  $\lambda \in \mathbb{T}_{>0}$ . Based on the two convexity notions we will also consider cones over the respective convex sets.

We will often identify a finite set with the columns of a matrix and vice versa.

**3.1. TO-convexity.** We collect several results from [42]. Note that TO-convexity appears under the name 'signed tropical convexity' there.

The *TO-convex hull* of a matrix  $A \in \mathbb{T}_\pm^{d \times n}$  is

$$\text{conv}_{\text{TO}}(A) = \bigcup \left\{ \mathcal{U}(A \odot x) \mid x \in \mathbb{T}_{\geq 0}^n, \bigoplus_{j \in [n]} x_j = 0 \right\} \subseteq \mathbb{T}_\pm^d. \quad (11)$$

A set  $M \subseteq \mathbb{T}_{\pm}^d$  is *TO-convex* if  $\text{conv}_{\text{TO}}(T) \subseteq M$  for all finite  $T \subseteq M$ . The hull operator extends by setting

$$\begin{aligned} \text{conv}_{\text{TO}}(M) &= \bigcap_{M \subseteq S, S \text{ TO-convex}} S \\ &= \bigcup_{T \subseteq M, T \text{ finite}} \text{conv}_{\text{TO}}(T) . \end{aligned}$$

For a vector  $s \in \mathbb{S}^d$  with possibly balanced entries, the set  $\mathcal{U}(s) \subseteq \mathbb{T}_{\pm}^d$  can be seen as a hypercube. If  $s$  has  $k$  entries in  $\mathbb{T}_{\bullet} \setminus \{\mathbf{0}\}$ , we say that the hypercube  $\mathcal{U}(s)$  has *dimension*  $k$ . Furthermore, we say that  $\mathcal{U}(s')$  is a *face* of  $\mathcal{U}(s)$  if  $s'_k = s_k$  for every  $k$  such that  $s_k \in \mathbb{T}_{\pm}$  and  $|s'_k| = |s_k|$  for every other  $k$ . A face of dimension zero is called a *vertex* of  $\mathcal{U}(s)$ . In particular, a hypercube of dimension  $k$  has  $2^k$  vertices.

We recall a crucial property of TO-convexity.

**Lemma 3.1** ([42, Proposition 3.6]). *A subset  $U \subseteq \mathbb{T}_{\pm}^d$  is TO-convex if and only if  $\text{conv}_{\text{TO}}(\{p, q\})$  is contained in  $U$  for all  $p, q \in U$ .*

The TO-convex hull of a finite set can also be given as intersection of open tropical halfspaces. This is the origin of the name ‘TO-convex’ derived from ‘tropical open’. Furthermore, it motivates the definition of the TC-convex hull (‘tropical closed’) in Section 3.2.

**Theorem 3.2** ([42, Theorem 5.1]). *The TO-convex hull of a finite subset  $M \subseteq \mathbb{T}_{\pm}^d$  equals the intersection of all open halfspaces containing  $M$ , i.e.,*

$$\text{conv}_{\text{TO}}(M) = \bigcap_{M \subseteq \mathcal{H}^+(a)} \mathcal{H}^+(a)$$

It turns out that TO-convexity has rather well-behaved separation properties: it fulfills the Pasch property and the Kakutani property. For this, we need the notion of a *TO-hemisphere*, this is a TO-convex subset  $X \subseteq \mathbb{T}_{\pm}^d$  for which also  $\mathbb{T}_{\pm}^d \setminus X$  is TO-convex. A similar statement was proven for the unsigned case in [35].

**Theorem 3.3.** *TO-convexity has the Pasch property and the Kakutani property. In other words,*

- (i) *(Pasch property) if  $(a, b_1, b_2, c_1, c_2) \in \mathbb{T}_{\pm}^d$  are such that  $b_1 \in \text{conv}_{\text{TO}}(a, c_1)$ ,  $b_2 \in \text{conv}_{\text{TO}}(a, c_2)$ , then  $\text{conv}_{\text{TO}}(c_1, b_2) \cap \text{conv}_{\text{TO}}(c_2, b_1) \neq \emptyset$ ;*
- (ii) *(Kakutani property) if  $A, B \subseteq \mathbb{T}_{\pm}^d$  are two disjoint TO-convex sets, then there exists a TO-hemisphere  $X \subseteq \mathbb{T}_{\pm}^d$  such that  $A \subseteq X$  and  $B \subseteq (\mathbb{T}_{\pm}^d \setminus X)$ .*

*Proof.* For (i), we follow the proof of the Pasch property over arbitrary ordered fields, see [51, I. Proposition 4.14.1]. Let  $(a, b_1, b_2, c_1, c_2)$  be as in (i). Then there are  $s_1, s_2, \bar{s}_1, \bar{s}_2 \in \mathbb{T}_{\geq 0}$  such that

$$s_1 \oplus \bar{s}_1 = 0 \quad , \quad s_2 \oplus \bar{s}_2 = 0 \tag{12}$$

and

$$b_1 \in \mathcal{U}(s_1 \odot a \oplus \bar{s}_1 \odot c_1) \quad \text{and} \quad b_2 \in \mathcal{U}(s_2 \odot a \oplus \bar{s}_2 \odot c_2) . \tag{13}$$




FIGURE 2. The Pasch property in the real plane.

We define

$$\begin{aligned}
t_1 &= s_2 \odot (s_1 \oplus \bar{s}_1 \odot s_2)^{\odot -1} \\
\bar{t}_1 &= s_1 \odot \bar{s}_2 \odot (s_1 \oplus \bar{s}_1 \odot s_2)^{\odot -1} \\
t_2 &= \bar{s}_1 \odot s_2 \odot (s_1 \oplus \bar{s}_1 \odot s_2)^{\odot -1} \\
\bar{t}_2 &= s_1 \odot (s_1 \oplus \bar{s}_1 \odot s_2)^{\odot -1} .
\end{aligned} \tag{14}$$

Using (12), we obtain

$$\begin{aligned}
s_2 \oplus s_1 \odot \bar{s}_2 &= s_1 \odot s_2 \oplus \bar{s}_1 \odot s_2 \oplus s_1 \odot \bar{s}_2 \\
&= s_1 \odot s_2 \oplus s_1 \odot \bar{s}_2 \oplus \bar{s}_1 \odot s_2 = s_1 \oplus \bar{s}_1 \odot s_2,
\end{aligned}$$

which implies

$$t_1 \oplus \bar{t}_1 = (s_2 \oplus s_1 \odot \bar{s}_2) \odot (s_1 \oplus \bar{s}_1 \odot s_2)^{\odot -1} = 0 .$$

Note that we also have  $t_2 \oplus \bar{t}_2 = 0$ . Hence, we obtain

$$\mathcal{U}(t_1 \odot b_1 \oplus \bar{t}_1 \odot c_2) \subseteq \text{conv}_{\text{TO}}(b_1, c_2) \quad \text{and} \quad \mathcal{U}(t_2 \odot c_1 \oplus \bar{t}_2 \odot b_2) \subseteq \text{conv}_{\text{TO}}(c_1, b_2) .$$

Multiplying by the denominator  $s_1 \oplus \bar{s}_1 \odot s_2$  in (14), we see that the intersection  $\text{conv}_{\text{TO}}(b_1, c_2) \cap \text{conv}_{\text{TO}}(c_1, b_2)$  is not empty if

$$\mathcal{U}(s_2 \odot b_1 \oplus s_1 \odot \bar{s}_2 \odot c_2) \cap \mathcal{U}(\bar{s}_1 \odot s_2 \odot c_1 \oplus s_1 \odot b_2) \neq \emptyset .$$

Scaling (13) to

$$s_2 \odot b_1 \in \mathcal{U}(s_2 \odot s_1 \odot a \oplus s_2 \odot \bar{s}_1 \odot c_1) \quad \text{and} \quad s_1 \odot b_2 \in \mathcal{U}(s_1 \odot s_2 \odot a \oplus s_1 \odot \bar{s}_2 \odot c_2)$$

allows to apply Corollary 2.8 by setting

$$v = s_1 \odot s_2 \odot a, q = s_2 \odot b_1, w = \bar{s}_1 \odot s_2 \odot c_1, p = s_1 \odot b_2, u = s_1 \odot \bar{s}_2 \odot c_2.$$

To prove (ii), we use [24, Theorem 5], which shows that the Pasch property and the Kakutani property are equivalent for 2-ary convexities (see also [41, Theorem 4.1] for a more recent improvement). TO-convexity is 2-ary by Lemma 3.1. This concludes the proof.  $\square$

*Example 3.4.* Let  $a = (0, \ominus 0)$ ,  $b_1 = (\mathbb{O}, \ominus 0)$ ,  $b_2 = (0, 0)$ ,  $c_1 = (\ominus 0, \ominus 0)$ , and  $c_2 = (\mathbb{O}, 1)$ . Figure 3 depicts this configuration of points. The TO-convex hull of  $a$  and  $c_1$  is the straight line connecting them, so  $b_1 \in \text{conv}_{\text{TO}}(a, c_1)$ . Furthermore,  $(-1) \odot c_2 \oplus a = (0, \bullet 0)$ , so  $b_2 \in \text{conv}_{\text{TO}}(a, c_2)$ . One gets that the TO-convex hull of  $b_1$  and  $c_2$  is the dashed line connecting them and the TO-convex hull of  $c_1$  and  $b_2$  is just the shaded whole square. Their non-empty intersection visualizes the Pasch property in this example.




FIGURE 3. The Pasch property in the TO-convexity is satisfied.




FIGURE 4. The Pasch property in the TC-convexity is not satisfied; see Example 3.16.

**3.2. TC-convexity.** While the structure of TO-convexity is heavily linked to open tropical halfspaces, we introduce TC-convexity based on closed tropical halfspaces. For technical reasons, we first only define the TC-convex hull of two points, and then extend it to arbitrary sets using these TC-convex line segments. Later it turns out that, for an arbitrary finite set, this is actually the same as just taking the intersection of all closed tropical halfspaces containing it.

For basics on general convexity, we refer to [51]. We will mainly rely on the notion of a *convexity structure* [51, §1.1]. In particular, we are interested in convexity structures induced by an *interval operator* [51, §4.1].

**Definition 3.5.** We define the *TC-convex hull* of two points  $x, y \in \mathbb{T}_{\pm}^d$  as

$$\text{conv}_{\text{TC}}(x, y) = \bigcap_{x, y \in \overline{\mathcal{H}^+(a)}} \overline{\mathcal{H}^+(a)}. \quad (15)$$

Sets of this form are *TC-convex intervals*.

**Definition 3.6.** We say that a set  $X \subset \mathbb{T}_{\pm}^d$  is *TC-convex* if  $\text{conv}_{\text{TC}}(x, y) \subseteq X$  for all  $x, y \in X$ .

We have the following desirable properties.

**Corollary 3.7.** *The TC-convex sets form a convexity structure. In particular, TC-convex sets are closed by intersection and nested union.*

The definition of TC-convexity directly gives two important classes of TC-convex sets.

**Corollary 3.8.** *A closed signed affine tropical halfspace and a signed affine tropical hyperplane is TC-convex. TC-convex intervals, i.e. sets of the form  $\text{conv}_{\text{TC}}(x, y)$ , for  $x, y \in \mathbb{T}_{\pm}^d$ , are TC-convex.*

*Proof.* Right from the definition we get that closed signed affine tropical halfspaces are TC-convex. The TC-convexity of the other sets follows from the intersection property.  $\square$

Corollary 3.7 allows us to extend the hull operator routinely to arbitrary sets  $M \subseteq \mathbb{T}_{\pm}^d$  by setting

$$\text{conv}_{\text{TC}}(M) = \bigcap_{M \subseteq S, S \text{ TC-convex}} S .$$

In particular, this convexity structure is ‘domain finite’ by [51, Theorem 1.3] which means the following.

**Corollary 3.9.** *For an arbitrary set  $M \subseteq \mathbb{T}_{\pm}^d$  we have*

$$\text{conv}_{\text{TC}}(M) = \bigcup_{T \subseteq M, T \text{ finite}} \text{conv}_{\text{TC}}(T) .$$

**Corollary 3.10.** *For an arbitrary set  $M \subseteq \mathbb{T}_{\pm}^d$  we have  $\text{conv}_{\text{TC}}(M) \subseteq \text{conv}_{\text{TO}}(M)$ . Furthermore, each TO-convex set is also TC-convex.*

*Proof.* For two point  $x, y \in \mathbb{T}_{\pm}^d$ , [42, Theorem 5.4] implies that  $\text{conv}_{\text{TO}}(x, y)$  is an intersection of finitely many closed halfspaces. Now, combining with the Definition 3.5 yields  $\text{conv}_{\text{TC}}(x, y) \subseteq \text{conv}_{\text{TO}}(x, y)$ . Therefore, by definition of a TC-convex set via TC-convex intervals (Definition 3.6), each TO-convex set is also TC-convex. Hence, using the extension of the hull operators for TO-convexity and TC-convexity to not necessarily finite sets gives the first claim.  $\square$




FIGURE 5. TO-convex intervals and TC-convex intervals in the plane (see Example 3.15)

To describe the TC-convex hull explicitly, we need significantly more tools than for the TO-convex hull.

**Definition 3.11.** Given a set of points  $X = \{x_1, \dots, x_n\} \in \mathbb{T}_\pm^d$ , we define

$$\text{Vert}(X) = \text{Vert}(x_1, \dots, x_n) := \{x_{\sigma(1)} \triangleleft x_{\sigma(2)} \triangleleft \dots \triangleleft x_{\sigma(n)} \mid \sigma \in \text{Sym}(n)\} \subset \mathbb{T}_\pm^d,$$

where  $\text{Sym}(n)$  denotes the group of permutations of  $[n]$ . Furthermore, we denote by

$$\text{Faces}(X) := \text{Faces}(x_1, \dots, x_n)$$

the union of all faces of  $\mathcal{U}(x_1 \oplus \dots \oplus x_n)$  (considering it as a hypercube) whose vertices belong to  $\text{Vert}(x_1, \dots, x_n)$ .

While the convex structure of cancellation for TO-convexity only depends on the balanced outcome, the non-associative structure of TC-convexity is far more subtle. Therefore, it is not enough to apply a univariate operator like  $\mathcal{U}(\cdot)$  but we have to use the multivariate operator  $\text{Faces}(\cdot)$ . Though, equipped with this tool, we will be able to describe the TC-hull also in terms of an analog of convex combinations.

Directly from the definition and Observation 2.13 we get the next.

**Corollary 3.12.** *The set  $\text{Vert}(x_1, \dots, x_n)$  is a subset of vertices of the hypercube  $\mathcal{U}(x_1 \oplus \dots \oplus x_n)$ . In particular, it contains at most  $2^d$  points.*

*Example 3.13.* If we only consider two points, there are essentially three cases. We illustrate them on small examples. In the first case, the sum does not have a balanced entry.

$$\text{Vert}\left(\begin{pmatrix} 0 \\ \ominus 0 \end{pmatrix}, \begin{pmatrix} 1 \\ -1 \end{pmatrix}\right) = \left\{\begin{pmatrix} 1 \\ \ominus 0 \end{pmatrix}\right\} = \text{Faces}\left(\begin{pmatrix} 0 \\ \ominus 0 \end{pmatrix}, \begin{pmatrix} 1 \\ -1 \end{pmatrix}\right).$$

In the second case, we have

$$\text{Vert}\left(\begin{pmatrix} 0 \\ \ominus 0 \end{pmatrix}, \begin{pmatrix} 1 \\ 0 \end{pmatrix}\right) = \left\{\begin{pmatrix} 1 \\ \ominus 0 \end{pmatrix}, \begin{pmatrix} 1 \\ 0 \end{pmatrix}\right\}$$

which yields

$$\text{Faces}\left(\begin{pmatrix} 0 \\ \ominus 0 \end{pmatrix}, \begin{pmatrix} 1 \\ 0 \end{pmatrix}\right) = \left\{\begin{pmatrix} 1 \\ s \end{pmatrix} : s \in [\ominus 0, 0]\right\}.$$

In the last case, there is more than one balanced entry

$$\text{Vert}\left(\begin{pmatrix} 0 \\ \ominus 0 \end{pmatrix}, \begin{pmatrix} \ominus 0 \\ 0 \end{pmatrix}\right) = \left\{\begin{pmatrix} 0 \\ \ominus 0 \end{pmatrix}, \begin{pmatrix} \ominus 0 \\ 0 \end{pmatrix}\right\} = \text{Faces}\left(\begin{pmatrix} 0 \\ \ominus 0 \end{pmatrix}, \begin{pmatrix} \ominus 0 \\ 0 \end{pmatrix}\right).$$

**Proposition 3.14.** *We have*

$$\text{conv}_{\text{TC}}(x, y) = \bigcup \{\text{Faces}(\lambda \odot x, \mu \odot y) \mid \lambda, \mu \in \mathbb{T}_{\geq 0}, \lambda \oplus \mu = 0\}. \quad (16)$$

*Proof.* By Corollary 3.10, we know that  $\text{conv}_{\text{TC}}(x, y) \subseteq \text{conv}_{\text{TO}}(x, y)$ . With (11), this implies that it suffices to consider which part of  $\mathcal{U}(\lambda \odot x \oplus \mu \odot y)$  is contained in  $\text{conv}_{\text{TC}}(x, y)$  for each pair  $\lambda, \mu \in \mathbb{T}_{\geq 0}, \lambda \oplus \mu = 0$ . Note that, for different such pairs  $\lambda_1, \mu_1$  and  $\lambda_2, \mu_2$ , the sets  $\mathcal{U}(\lambda_1 \odot x \oplus \mu_1 \odot y)$  and  $\mathcal{U}(\lambda_2 \odot x \oplus \mu_2 \odot y)$  are either disjoint or identical. Therefore, it is enough to consider the sets  $\mathcal{U}(\lambda \odot x \oplus \mu \odot y)$  for a fixed pair  $\lambda, \mu$ . Hence, fixing such a pair, we distinguish three cases.

For  $a = (a_0, \bar{a}) \in \mathbb{T}_\pm^{d+1}$  let  $\overline{\mathcal{H}}^+(a)$  be an affine halfspace containing  $x$  and  $y$ . This means that

$$\lambda \odot a_0 \oplus \bar{a} \odot \lambda \odot x \neq \mathbf{0} \text{ and } \mu \odot a_0 \oplus \bar{a} \odot \mu \odot y \neq \mathbf{0}. \quad (17)$$

Recall that at least one of  $\lambda, \mu$  is 0.

**Case 1** ( $\lambda \odot x \oplus \mu \odot y$  has no balanced entry. ) Adding the relations in (17) yields

$$a_0 \oplus \bar{a} \odot (\lambda \odot x \oplus \mu \odot y) \vDash \mathbf{0} .$$

That already concludes this case, where indeed  $\lambda \odot x \oplus \mu \odot y = \lambda \odot x \triangleleft \mu \odot y = \mu \odot y \triangleleft \lambda \odot x$ .

**Case 2** ( $\lambda \odot x \oplus \mu \odot y$  has exactly one balanced entry. ) Without loss of generality, we assume that the  $d$ -th entry is balanced. Reformulating (17) implies

$$\begin{aligned} \lambda \odot a_0 \oplus \bigoplus_{\ell=1}^{d-1} a_\ell \odot \lambda \odot x_\ell &\vDash \ominus \lambda \odot a_d \odot x_d \\ \mu \odot a_0 \oplus \bigoplus_{\ell=1}^{d-1} a_\ell \odot \mu \odot y_\ell &\vDash \ominus \mu \odot a_d \odot y_d . \end{aligned}$$

Using  $b := \lambda \odot x_d = \ominus \mu \odot y_d$ , the side-wise addition of the relations yields

$$a_0 \oplus \bigoplus_{\ell=1}^{d-1} a_\ell \odot (\lambda \odot x_\ell \oplus \mu \odot y_\ell) \vDash a_d \oplus \bullet b .$$

Therefore, each point in

$$\{(\lambda \odot x_1 \oplus \mu \odot y_1, \dots, \lambda \odot x_{d-1} \oplus \mu \odot y_{d-1}, s) : s \in [\ominus|b|, |b|]\}$$

is contained as claimed. Furthermore, the latter set is indeed the line between  $\lambda \odot x \triangleleft \mu \odot y$  and  $\mu \odot y \triangleleft \lambda \odot x$ .

**Case 3** ( $\lambda \odot x \oplus \mu \odot y$  has more than one balanced entry. ) Without loss of generality, exactly the coordinates  $k+1$  to  $d$  of  $\lambda \odot x \oplus \mu \odot y$  are balanced. Then (17) amounts to

$$\begin{aligned} \left( \lambda \odot a_0 \oplus \bigoplus_{\ell=1}^k a_\ell \odot \lambda \odot x_\ell \right) \oplus \left( \bigoplus_{\ell=k+1}^d a_\ell \odot \lambda \odot x_\ell \right) &\vDash \mathbf{0} \\ \left( \mu \odot a_0 \oplus \bigoplus_{\ell=1}^k a_\ell \odot \mu \odot y_\ell \right) \oplus \left( \bigoplus_{\ell=k+1}^d a_\ell \odot \mu \odot y_\ell \right) &\vDash \mathbf{0} . \end{aligned}$$

Using Lemma 2.6 in the same way as we did for Case 2, we obtain that  $\lambda \odot x \triangleleft \mu \odot y$  and  $\mu \odot y \triangleleft \lambda \odot x$  are also contained in  $\overline{\mathcal{H}}^+(a)$ .

Up until now, we have proven the first inclusion. Now, we set  $b_\ell = \lambda \odot x_\ell = \ominus \mu \odot y_\ell$  for  $\ell \in [d] \setminus [k]$ . We look at the halfspaces

$$\overline{\mathcal{H}}^+(\ominus 0, \mathbf{0}, \dots, \mathbf{0}, \epsilon_1 \odot b_{k+1}^{\odot -1}, \dots, \epsilon_{d-k} \odot b_d^{\odot -1}) \quad (18)$$

for each  $\epsilon \in \{\ominus, \oplus\}^{d-k} \setminus \{(\ominus, \dots, \ominus), (\oplus, \dots, \oplus)\}$ . By putting  $x$  and  $y$  in the corresponding relation, we see that they are both contained.

Now let  $w_1 = \lambda \odot x \triangleleft \mu \odot y$  and  $w_2 = \mu \odot y \triangleleft \lambda \odot x$ . Note that  $w_1$  and  $x$  have the same sign pattern on the coordinates  $k+1$  up to  $d$ , and the same for  $w_2$  and  $y$ , respectively. We pick any point

$$z \in \mathcal{U}(\lambda \odot x_1 \oplus \mu \odot y_1, \dots, \lambda \odot x_k \oplus \mu \odot y_k, \bullet b_{k+1}, \dots, \bullet b_d) \setminus \{w_1, w_2\}$$

**Case 3a** ( $z$  has the same sign pattern as  $w_1$  or  $w_2$ . ) Without loss of generality, we assume that  $z$  has the same sign pattern as  $w_1$ . Then there is a coordinate of

$z$ , say the  $(k+1)$ st, such that  $|z_{k+1}| < |\lambda \odot x_{k+1}| = |\mu \odot y_{k+1}|$ . We let

$$(\epsilon_1, \dots, \epsilon_{d-k}) = (\text{tsgn}(z_{k+1}), \ominus \text{tsgn}(z_{k+2}), \dots, \ominus \text{tsgn}(z_{d-k}))$$

As  $|z_{k+1} \odot |b_{k+1}|^{\odot -1}| < 0$ , we get

$$\ominus 0 \oplus \epsilon_1 \odot |b_{k+1}|^{\odot -1} \odot z_{k+1} \oplus \dots \oplus \epsilon_{d-k} \odot |b_d|^{\odot -1} \odot z_{d-k} < \mathbf{0}.$$

Hence, there is a halfspace among those in (18) not containing  $z$ .

**Case 3b** ( $z$  has a different sign pattern from  $w_1$  and  $w_2$ .) We let

$$(\epsilon_1, \dots, \epsilon_{d-k}) = (\ominus \text{tsgn}(z_{k+1}), \dots, \ominus \text{tsgn}(z_{d-k}))$$

be the negative of the sequence of signs. Because of the relation

$$\ominus 0 \oplus \epsilon_1 \odot |b_{k+1}|^{\odot -1} \odot z_{k+1} \oplus \dots \oplus \epsilon_{d-k} \odot |b_d|^{\odot -1} \odot z_{d-k} < \mathbf{0}$$

there is a halfspace among those in (18) not containing  $z$ , as these include all sign patterns except for those of  $x$  and  $y$ .  $\square$

*Example 3.15.* Figure 5 compares TC-intervals with TO-intervals in a plane. If we take  $x = (0, 0)$  and  $y = (\ominus - 3, \ominus - 2)$ , then for every  $\lambda, \mu \in \mathbb{T}_{\geq \mathbf{0}}$  such that  $\lambda \oplus \mu = 0$  we have  $\text{Faces}(\lambda \odot x, \mu \odot y) = \mathcal{U}(\lambda \odot x \oplus \mu \odot y)$ . In particular, in this case the intervals coincide,  $\text{conv}_{\text{TO}}(x, y) = \text{conv}_{\text{TC}}(x, y)$ . If we take  $y = (\ominus - 2, \ominus - 2)$  instead, then the equality no longer holds. Indeed, for  $\lambda = -2$ ,  $\mu = 0$  we have  $\mathcal{U}(\lambda \odot x \oplus \mu \odot y) = \mathcal{U}(\bullet 2, \bullet 2) = [\ominus - 2, -2] \times [\ominus - 2, -2]$  and  $\text{Faces}(\lambda \odot x, \mu \odot y) = \{(\ominus - 2, \ominus - 2), (-2, -2)\}$ .

The next example shows that the TC-convexity does not have the Pasch property or the Kakutani property. Nevertheless, in Section 5 we will show that the TC-convexity satisfies a weaker separation property.

*Example 3.16.* As in Example 3.4, let  $a = (0, \ominus 0)$ ,  $b_1 = (\mathbf{0}, \ominus 0)$ ,  $b_2 = (0, 0)$ ,  $c_1 = (\ominus 0, \ominus 0)$ , and  $c_2 = (\mathbf{0}, 1)$ . Figure 4 depicts this configuration of points. Note that we have  $b_1 \in \text{Faces}(a, c_1)$  and  $b_2 \in \text{Faces}(a, -1 \odot c_2)$ , so that  $b_1 \in \text{conv}_{\text{TC}}(a, c_1)$ ,  $b_2 \in \text{conv}_{\text{TC}}(a, c_2)$ . Nevertheless, the interval  $\text{conv}_{\text{TC}}(c_1, b_2)$  is reduced to two points,  $\text{conv}_{\text{TC}}(c_1, b_2) = \{c_1, b_2\}$ , while  $\text{conv}_{\text{TC}}(c_2, b_1) = \{\mathbf{0}\} \times [\ominus 0, 1]$ . In particular, the intersection  $\text{conv}_{\text{TC}}(c_1, b_2) \cap \text{conv}_{\text{TC}}(c_2, b_1)$  is empty, so the TC-convexity does not have the Pasch property. This also shows that the TC-convexity does not have the Kakutani property: there is no TC-hemisphere that separates  $\text{conv}_{\text{TC}}(c_1, b_2)$  from  $\text{conv}_{\text{TC}}(c_2, b_1)$ , because the point  $a$  could not belong to either side of this hemisphere. Even further, we note that  $\text{conv}_{\text{TC}}(b_1, c_2) = \text{conv}_{\text{TO}}(b_1, c_2)$ . Hence, this example shows that the TO-convexity and TC-convexity do not satisfy the Kakutani property for pairs of convexities studied in [41].

**Proposition 3.17.** *A set  $M \subseteq \mathbb{T}_{\pm}^d$  is TC-convex if and only if it is closed under the following two operations:*

(i) *for  $x, y \in M$  and  $\lambda, \mu \in \mathbb{T}_{\geq \mathbf{0}}$  with  $\lambda \oplus \mu = 0$ , we have*

$$\lambda \odot x \triangleleft \mu \odot y \in M; \quad (\text{weighted left sum})$$

(ii) *if  $(u, v, w), (u, \ominus v, w) \in \mathbb{T}_{\pm}^k \times \mathbb{T}_{\pm} \times \mathbb{T}_{\pm}^{d-k-1}$  for  $k \in [d]_0$  are contained in  $M$  then*

$$\{u\} \times [\ominus|v|, |v|] \times \{w\} \subseteq M. \quad (\text{local elimination})$$



*Proof.* By Proposition 3.14, a TC-convex set fulfills these two properties.

It suffices to prove that the TC-convex hull of two points can be generated by these two operations. So we fix two points  $x, y \in M \subseteq \mathbb{T}_{\pm}^d$ . Let  $\lambda, \mu \in \mathbb{T}_{\geq 0}$  with  $\lambda \oplus \mu = 0$ . By the first property,  $\text{Vert}(\lambda \odot x, \mu \odot y)$  is contained in  $M$ . If  $\text{Vert}(\lambda \odot x, \mu \odot y)$  do not form the vertices of a face of the respective hypercube, it equals  $\text{Faces}(\lambda \odot x, \mu \odot y)$ . Otherwise  $\text{Faces}(\lambda \odot x, \mu \odot y)$  is the line segment arising from the second property. Now, Proposition 3.14 concludes the proof.  $\square$

*Example 3.18.* For  $(p, r), (q, r) \in \mathbb{T}_{\pm} \times \mathbb{T}_{\pm}^{d-1}$  with  $p < q$ , one gets

$$\text{conv}_{\text{TC}}(\{(p, r), (q, r)\}) = [p, q] \times \{r\}. \quad (19)$$

To see this, we first assume that  $p < q$  have the same sign, w.l.o.g. both are positive. Then  $[p, q] \times \{r\} = \{(p, r) \triangleleft \lambda \odot (q, r) \mid 0 \leq \lambda \leq 0\}$ . Otherwise, assume that  $p < 0$  and  $|p| < q$ . Then  $(\ominus p, r) = (q - |p|) \odot (q, r) \triangleleft (p, r)$  and the claim follows with the local elimination property. The other cases follow by suitably adapting the signs and scalars.

For  $p, q \in \mathbb{T}_{\pm}^d$  with  $p \leq q$  (defined component-wise), we denote

$$\begin{aligned} [p, q] &= \{x \in \mathbb{T}_{\pm}^d \mid p_i \leq x_i \leq q_i \forall i \in [d]\}, \\ (p, q) &= \{x \in \mathbb{T}_{\pm}^d \mid p_i < x_i < q_i \forall i \in [d]\}. \end{aligned}$$

*Example 3.19.* For  $p, q \in \mathbb{T}_{\pm}^d$  with  $p \leq q$ , one gets

$$\text{conv}_{\text{TC}}(\{p_1, q_1\} \times \cdots \times \{p_d, q_d\}) = [p, q]. \quad (20)$$

To see this, we can fix all but one coordinate and then iteratively use Example 3.18.

Note that it also holds that

$$\text{conv}_{\text{TO}}(\{p_1, q_1\} \times \cdots \times \{p_d, q_d\}) = [p, q]. \quad (21)$$

Let  $C$  be a subcomplex of the faces of  $[-1, 1]^d$  with the property: *if all vertices of a face are contained in  $C$  then so is the face.*

For  $\sim \in \{\leq, \geq\}^d$ , we define  $x \sim y \Leftrightarrow (x_k \sim_k y_k \text{ for all } k \in [d])$ .

**Lemma 3.20.** *A point  $z \in [-1, 1]^d$  is contained in  $C$  if and only if for every  $\sim \in \{\leq, \geq\}^d$  there exists a vertex  $w$  of  $C$  such that  $z \sim w$ .*

*Proof.* With a point  $z \in [-1, 1]^d$ , we associate a partition  $I \cup J \cup K$  of  $[d]$  indexing the coordinates of  $z$  which are  $+1$ ,  $-1$  or in the open interval  $(-1, 1)$ . Then the set  $I \cup J$  uniquely defines the face  $F$  of the cube of smallest dimension containing  $z$ .

Assume that  $z$  is not contained in  $C$ . By the crucial property of  $C$ , there is a vertex  $v$  of  $F$  which is not contained in  $C$ . With  $v$  we associate a vector  $\sim^{(v)} \in \{\leq, \geq\}^d$  by converting  $-1$  to  $\leq$  and  $+1$  to  $\geq$ . Then  $v$  is the unique vertex of  $[-1, 1]^d$  with  $v \sim^{(v)} z$  because there is a unique choice of  $\tau \in \{-1, 1\}$  with  $\tau \sim_k^{(v)} z_k$  for  $k \in K$ . As  $v$  is not a vertex of  $C$  this concludes the first direction.

On the other hand, if a point  $z$  is contained in  $C$  then also the face  $F$  is contained in  $C$ . Let  $\sim \in \{\leq, \geq\}^d$  be arbitrary. We define a vertex  $v$  of  $[-1, 1]^d$  by  $v_\ell = z_\ell$  for  $\ell \in I \cup J$  and

$$v_k = \begin{cases} 1 & \text{if } \sim_k \text{ equals } \geq, \\ -1 & \text{if } \sim_k \text{ equals } \leq \end{cases}$$

for  $k \in K$ . By construction,  $v$  is a vertex of  $F$  and, hence, of  $C$ . Furthermore, we get  $v \sim z$  proving the claim.  $\square$

**Corollary 3.21.** *We have  $y \in \text{Faces}(x_1, \dots, x_n)$  if and only if for every  $\sim \in \{\leq, \geq\}^d$  there exists  $w \in \text{Vert}(x_1, \dots, x_n)$  such that  $y_k \sim_k w_k$  for all  $k \in [d]$ .*

*Proof.* By Definition 3.11, the set  $\text{Faces}(x_1, \dots, x_n)$  forms a subcomplex of the faces of the (stretched) hypercube  $\mathcal{U}(x_1 \oplus \dots \oplus x_n)$  exactly with the property required of  $C$  in Lemma 3.20. As  $\text{Vert}(x_1, \dots, x_n)$  select exactly the vertices of this subcomplex, Lemma 3.20 gives the required equivalence.  $\square$

**3.3. Fundamental properties of TC-convex sets.** Fix finite sets  $X, Y \subseteq \mathbb{T}_{\pm}^d$ .

**Lemma 3.22.** *Let  $s, t \in \mathbb{S}$  with  $|s| = |\bigoplus X|$  and  $|t| = |\bigoplus Y|$  such that  $\mathcal{U}(s) \subseteq \text{Faces}(X)$  and  $\mathcal{U}(t) \subseteq \text{Faces}(Y)$ . Then  $\mathcal{U}(s \triangleleft t) \subseteq \text{Faces}(X \cup Y)$ .*

*Proof.* Note that, by definition of  $s$ , we have the inclusion  $\mathcal{U}(\bullet s) \subseteq \mathcal{U}(\bigoplus X)$  and further, using  $\mathcal{U}(s) \subseteq \text{Faces}(X)$ , that every vertex of  $\mathcal{U}(s)$  is a vertex of  $\mathcal{U}(\bigoplus X)$ . Analogously, every vertex of  $\mathcal{U}(t)$  is a vertex of  $\mathcal{U}(\bigoplus Y)$ .

Therefore as part of  $\text{Faces}(\bigoplus X)$ , all vertices of  $\mathcal{U}(s)$  are of the form  $\triangleleft X^\sigma$  for some ordering  $\sigma$  of  $X$ , and analogously, all vertices of  $\mathcal{U}(t)$  are of the form  $\triangleleft Y^\tau$  for some ordering  $\tau$  of  $Y$ . Now, we consider a vertex  $u$  of  $\mathcal{U}(s \triangleleft t)$ . We partition  $[d]$  in two sets,  $I := \{i: |u_i| = |s_i|\}$  and its complement.

By definition of the left sum,  $u$  is given by (signed versions of) the entries of  $s$  on  $I$  and (signed versions of) the entries of  $t$  on  $[d] \setminus I$ . Let  $v$  be a vertex of  $\mathcal{U}(s)$  which agrees with  $u$  on  $I$  and let  $w$  be a vertex of  $\mathcal{U}(t)$  which agrees with  $u$  on  $[d] \setminus I$ . This just means that  $u = v \triangleleft w$ . Furthermore, by construction, the latter left sum lies in  $\text{Vert}(X \cup Y)$ . As  $u$  belongs to  $\text{Vert}(X \cup Y)$ , we get the inclusion  $\mathcal{U}(s \triangleleft t) \subseteq \text{Faces}(X \cup Y)$  by definition of  $\text{Faces}(\cdot)$ .  $\square$

**Lemma 3.23.** *The set  $\text{Faces}(X)$  is TC-convex.*

*In particular, we have  $\text{Faces}(X) = \text{conv}_{\text{TC}}(\text{Vert}(X))$*

*Proof.* Let  $x = |\bigoplus X|$  and let  $x^\sigma$  for  $\sigma \in \{\ominus, \oplus\}^d$  denote a signed version of  $x$ . Recall that the points  $\text{Vert}(X)$  are of the form  $x^\sigma$  where  $\sigma$  ranges over a subset  $\Sigma$  of  $\{\ominus, \oplus\}^d$ . Let  $H_\tau^+ = \overline{\mathcal{H}}^+(0, (x_1^\tau)^{\ominus-1}, \dots, (x_d^\tau)^{\ominus-1})$  be the halfspace with its ‘apex’ at the point  $x^\tau$  for  $\tau \in \{\ominus, \oplus\}^d$  containing  $\mathbb{O}$  and  $H_\tau^-$  the opposite closed one. Then the intersection  $\bigcap_{\tau \in \{\ominus, \oplus\}^d} H_\tau^+$  exactly yields the hypercube  $\mathcal{U}(\bullet x)$ . If we further intersect this with all halfspaces  $H_\tau^-$  for  $\tau \in \{\ominus, \oplus\}^d \setminus \Sigma$  we get  $\text{Faces}(X)$  as  $H_\tau^-$  exactly cuts off all faces of the cube containing  $x^\tau$  with  $\tau$  not in  $\Sigma$ . Hence, it is TC-convex as an intersection of TC-convex sets.

Now, we look at a face  $F$  of  $\mathcal{U}(\bullet x)$  for which all vertices  $V$  are contained in  $\text{Vert}(X)$ . Applying Proposition 3.14 iteratively on pairs of points, which only differ in the sign of one component, we get  $\text{Faces}(V) = F \subseteq \text{conv}_{\text{TC}}(V)$ . Ranging over all faces in  $\text{Faces}(X)$  yields  $\text{Faces}(X) \subseteq \text{conv}_{\text{TC}}(X)$ . But as  $\text{Faces}(X)$  is TC-convex, we get an equality.  $\square$

**Proposition 3.24.** *Let  $X \subseteq \mathbb{T}_{\pm}^d$  be an  $n$ -element set interpreted as a matrix. Then*

$$\text{conv}_{\text{TC}}(X) = \bigcup \left\{ \text{Faces}(X \odot \text{diag}(\lambda)) \mid \lambda \in \mathbb{T}_{\geq 0}^n, \bigoplus_i \lambda_i = 0 \right\}. \quad (22)$$

*Proof.* Let  $x_1, \dots, x_n \in X$  and  $\lambda_1, \dots, \lambda_n \in \mathbb{T}_{\geq 0}$  be such that  $\bigoplus_i \lambda_i = 0$ . We will start by showing that  $y = \lambda_1 \odot x_1 \triangleleft \dots \triangleleft \lambda_n \odot x_n$  belongs to  $\text{conv}_{\text{TC}}(X)$ . The proof goes by induction over  $n$ .

The claim is trivial for  $n = 1$ . For higher  $n$ , let  $\mu = \bigoplus_{i>1} \lambda_i$ . The claim is trivial if  $\mu = \mathbf{0}$ . Otherwise, let  $\mu_i = \lambda_i - \mu$  for every  $i > 1$ . Then, we have  $\lambda_1 \odot x_1 \triangleleft \dots \triangleleft \lambda_n \odot x_n = \lambda_1 \odot x_1 \triangleleft \mu \odot (\mu_2 \odot x_2 \triangleleft \dots \triangleleft \mu_n \odot x_n)$ . By the induction hypothesis, the point  $z = \mu_2 \odot x_2 \triangleleft \dots \triangleleft \mu_n \odot x_n$  belongs to  $\text{conv}_{\text{TC}}(X)$ . Therefore, the point  $y = \lambda_1 \odot x_1 \triangleleft \mu \odot z$  belongs to  $\text{conv}_{\text{TC}}(X)$  by Proposition 3.14. Thus, we have proven that every point of the form  $\lambda_1 \odot x_1 \triangleleft \dots \triangleleft \lambda_n \odot x_n$  belongs to  $\text{conv}_{\text{TC}}(X)$ . In particular, the set  $\text{Vert}(X \odot \text{diag}(\lambda))$  belongs to  $\text{conv}_{\text{TC}}(X)$ . Hence, Lemma 3.23 implies that  $\text{Faces}(X \odot \text{diag}(\lambda)) = \text{conv}_{\text{TC}}(\text{Vert}(X \odot \text{diag}(\lambda)))$  is contained in  $\text{conv}_{\text{TC}}(X)$ .

To finish the proof, we will show that the set on the right-hand side of (22) is TC-convex. To do so, suppose that  $a \in \text{Faces}(X \odot \text{diag}(\lambda))$  and  $b \in \text{Faces}(X \odot \text{diag}(\mu))$  for some  $\lambda_i, \mu_i \in \mathbb{T}_{\geq 0}$  and  $\bigoplus_i \lambda_i = \bigoplus_i \mu_i = 0$ . Further, let  $\alpha, \beta \in \mathbb{T}_{\geq 0}$  be such that  $\alpha \oplus \beta = 0$ . By Lemma 3.22, the set  $\text{Vert}(\alpha \odot a, \beta \odot b)$  is included in  $\text{Faces}(\alpha \odot X \odot \text{diag}(\lambda) \cup \beta \odot X \odot \text{diag}(\mu))$ . Hence,  $\text{Faces}(\alpha \odot a, \beta \odot b)$  is also included in this set by Lemma 3.23. Thus, by Proposition 3.14,  $\text{conv}_{\text{TC}}(a, b)$  is included in the set defined on the right-hand side of (22) since  $\text{Faces}(\alpha \odot X \odot \text{diag}(\lambda) \cup \beta \odot X \odot \text{diag}(\mu)) = \text{Faces}(X \odot \text{diag}(\nu))$  with  $\nu = \alpha \odot \text{diag}(\lambda) \oplus \beta \odot \text{diag}(\mu)$ , which fulfills  $\nu \in \mathbb{T}_{\geq 0}^n$  and  $\bigoplus_i \nu_i = 0$ .  $\square$

Using the representation of the convex hull as union of finite convex hulls stated in Corollary 3.9 we get the following.

**Corollary 3.25.** *If  $X \subseteq \mathbb{T}_{\pm}^d$ , then*

$$\text{conv}_{\text{TC}}(X) = \bigcup \left\{ \text{Faces}(X \odot \text{diag}(\lambda)) \mid \lambda \in \mathbb{T}_{\geq 0}^X, \bigoplus_i \lambda_i = 0, |\text{supp}(\lambda)| < +\infty \right\}.$$

We now give estimates on the Carathéodory-number of TC-convexity. The core case for this is the representation of vertices from a small set of generators.

**Lemma 3.26.** *Let  $X = \{x_1, \dots, x_n\} \subseteq \mathbb{T}_{\pm}^d$ . Then there is a subset  $Y \subseteq X$  with  $|Y| \leq d2^d$  and  $\text{Vert}(X) \subseteq \text{Vert}(Y)$ , hence,  $\text{Vert}(X) = \text{Vert}(Y)$ .*

*Proof.* Let  $y \in \text{Vert}(X)$  and let  $\sigma \in \text{Sym}(n)$  such that  $y = x_{\sigma(1)} \triangleleft \dots \triangleleft x_{\sigma(n)}$ . For every coordinate  $k \in [d]$  let  $j_k \in [n]$  be the smallest number such that  $y_k = (x_{\sigma(j_k)})_k$  and let  $J_y = \{\sigma(j_k) \mid k \in [d]\} \subseteq [n]$ . Note that  $|J_y| \leq d$ . Now, let  $I_y = (i_1, \dots, i_{|J_y|})$  be the ascending sequence of the elements in  $J_y$ . Observe that for any set  $J$  with  $J_y \subseteq J \subseteq [n]$ , it holds  $y = x_{i_1} \triangleleft \dots \triangleleft x_{|J_y|} \triangleleft \triangleleft_{i \in J \setminus J_y} x_i$  for any order of the summands indexed by  $J \setminus J_y$ .

Finally, we define  $J$  to be the union  $\bigcup_{y \in \text{Vert}(X)} J_y$ . By the above reasoning, we have  $|J| \leq d2^d$  and  $\text{Vert}(X) \subseteq \text{Vert}(Y)$ . As each left-sum of the elements in  $Y$  already defines a point for which the absolute values of the components equal the components of  $\bullet \bigoplus X$ , we also get the reverse inclusion  $\text{Vert}(Y) \subseteq \text{Vert}(X)$ .  $\square$

**Proposition 3.27.** *If  $X \subseteq \mathbb{T}_{\pm}^d$ , then*

$$\text{conv}_{\text{TC}}(X) = \bigcup \left\{ \text{Faces}(X \odot \text{diag}(\lambda)) \mid \lambda \in \mathbb{T}_{\geq 0}^X, \bigoplus_i \lambda_i = 0, |\text{supp}(\lambda)| \leq c_d \right\},$$

where  $c_d = d2^d + 1$ .

*Proof.* Let  $y \in \text{conv}_{\text{TC}}(X)$ . By Corollary 3.25 there is some  $\lambda \in \mathbb{T}_{\geq \mathbf{0}}^X$  with  $\bigoplus_i \lambda_i = 0$  such that  $y \in \text{Faces}(X \odot \text{diag}(\lambda))$ .

Let  $j_0$  be an index with  $\lambda_{j_0} = 0$ . We derive a new coefficient vector  $\mu$  from  $\lambda$  with  $\mu_{j_0} = 0$ ,  $\text{Vert}(X \odot \text{diag}(\lambda)) = \text{Vert}(X \odot \text{diag}(\mu))$  and  $|\text{supp}(\mu)| \leq c_d = d2^d + 1$ . Lemma 3.26 implies that we can achieve this by setting all but  $c_d$  entries of  $\lambda$  to  $\mathbf{0}$ .

With Lemma 3.23, we obtain

$$\begin{aligned} y \in \text{Faces}(X \odot \text{diag}(\lambda)) &= \text{conv}_{\text{TC}}(\text{Vert}(X \odot \text{diag}(\lambda))) \\ &\subseteq \text{conv}_{\text{TC}}(\text{Vert}(X \odot \text{diag}(\mu))) \\ &= \text{Faces}(X \odot \text{diag}(\mu)). \end{aligned}$$

Taking the subset  $Y$  of  $X$  given by the support of  $\mu$ , one sees  $y \in \text{conv}_{\text{TC}}(Y)$ .  $\square$

While TC-convexity extends the ‘usual’ tropical convexity whose Carathéodory number is  $d + 1$  as discussed in [32], we already get an exponential lower bound from a simple example.

*Example 3.28.* Let  $X = \{\ominus 0, 0\}^d$ . Then, Lemma 3.23 shows that  $\text{conv}_{\text{TC}}(X) = [\ominus 0, 0]^d$ . In particular, we have  $\mathbf{0} \in \text{conv}_{\text{TC}}(X)$ . Moreover, if  $Y \subsetneq X$  is any strict subset of  $X$ , then Lemma 3.23 shows that  $\text{conv}_{\text{TC}}(Y)$  is the union of faces of  $[\ominus 0, 0]^d$  whose vertices belong to  $Y$ . In particular,  $\mathbf{0} \notin \text{conv}_{\text{TC}}(Y)$ . This example shows the lower bound  $c_d \geq 2^d$  for the Carathéodory number of TC-convexity. We do not know what is the optimal value of  $c_d$ .

**Definition 3.29.** Given an arbitrary set  $X \subseteq \mathbb{T}_{\pm}^d$ , we denote

$$\text{cone}_{\text{TC}}(X) = \{ \lambda \odot x \mid \lambda \in \mathbb{T}_{\geq \mathbf{0}}, x \in \text{conv}_{\text{TC}}(X) \}.$$

**Lemma 3.30.** *If  $X \subseteq \mathbb{T}_{\pm}^d$ , then*

$$\text{cone}_{\text{TC}}(X) = \bigcup \{ \text{Faces}(X \odot \text{diag}(\lambda)) \mid \lambda \in \mathbb{T}_{\geq \mathbf{0}}^X, |\text{supp}(\lambda)| < +\infty \}.$$

*Proof.* By Proposition 3.24 we have the equality

$$\begin{aligned} &\text{cone}_{\text{TC}}(X) \\ &= \bigcup \left\{ \mu \odot \text{Faces}(X \odot \text{diag}(\lambda)) \mid \mu \in \mathbb{T}_{\geq \mathbf{0}}, \lambda \in \mathbb{T}_{\geq \mathbf{0}}^X, \bigoplus_i \lambda_i = 0, |\text{supp}(\lambda)| < +\infty \right\} \\ &= \bigcup \left\{ \text{Faces}(X \odot \text{diag}(\mu \odot \lambda)) \mid \mu \in \mathbb{T}_{\geq \mathbf{0}}, \lambda \in \mathbb{T}_{\geq \mathbf{0}}^X, \bigoplus_i \lambda_i = 0, |\text{supp}(\lambda)| < +\infty \right\} \end{aligned}$$

Furthermore, if  $\xi \in \mathbb{T}_{\geq \mathbf{0}}^X$ ,  $\xi \neq \mathbf{0}$  has finite support, then we can write it as  $\xi = \mu \odot \lambda$ , where  $\mu = \xi_1 \oplus \dots \oplus \xi_d \in \mathbb{T}_{\geq \mathbf{0}}$  and  $\lambda \in \mathbb{T}_{\geq \mathbf{0}}^X$ ,  $\bigoplus_i \lambda_i = 0$  is defined as  $\lambda_i = \xi_i \odot \mu^{\ominus -1}$  for all  $i$ . Therefore, we get

$$\begin{aligned} &\text{cone}_{\text{TC}}(X) \\ &= \bigcup \left\{ \text{Faces}(X \odot \text{diag}(\mu \odot \lambda)) \mid \mu \in \mathbb{T}_{\geq \mathbf{0}}, \lambda \in \mathbb{T}_{\geq \mathbf{0}}^X, \bigoplus_i \lambda_i = 0, |\text{supp}(\lambda)| < +\infty \right\} \\ &= \bigcup \{ \text{Faces}(X \odot \text{diag}(\lambda)) \mid \lambda \in \mathbb{T}_{\geq \mathbf{0}}^X, |\text{supp}(\lambda)| < +\infty \}. \quad \square \end{aligned}$$

**Corollary 3.31.** *The set  $\text{cone}_{\text{TC}}(X)$  is the smallest TC-convex cone that contains  $X$  and  $\mathbf{0}$ .*

*Proof.* We set  $\tilde{X} = X \cup \mathbf{0}$  and use the representation from Lemma 3.30 to get

$$\begin{aligned} & \text{cone}_{\text{TC}}(X) \\ &= \bigcup_{\mu \in \mathbb{T}_{\geq \mathbf{0}}} \bigcup \left\{ \text{Faces}(X \odot \text{diag}(\lambda)) \mid \lambda \in \mathbb{T}_{\geq \mathbf{0}}^X, \bigoplus_i \lambda_i \leq \mu, |\text{supp}(\lambda)| < +\infty \right\} \\ &= \bigcup_{\mu \in \mathbb{T}_{\geq \mathbf{0}}} \bigcup \left\{ \text{Faces}(\tilde{X} \odot \text{diag}(\lambda)) \mid \lambda \in \mathbb{T}_{\geq \mathbf{0}}^{\tilde{X}}, \bigoplus_i \lambda_i = \mu, |\text{supp}(\lambda)| < +\infty \right\} \\ &= \bigcup_{\mu \in \mathbb{T}_{\geq \mathbf{0}}} \text{conv}_{\text{TC}}(\mu \odot \tilde{X}) . \end{aligned}$$

As this is a nested union for increasing  $\mu$ , it is a TC-convex set by Corollary 3.7.

Furthermore, let  $Z$  be the smallest TC-convex cone containing  $\tilde{X}$ . By definition of a cone, one gets  $\mu \odot \tilde{X} \subseteq Z$  for all  $\mu > \mathbf{0}$  and, by TC-convexity, also  $\text{conv}_{\text{TC}}(\mu \odot \tilde{X}) \subseteq Z$  for all  $\mu > \mathbf{0}$ . Since  $\mathbf{0}$  belongs to  $Z$ , the equality above shows the minimality of  $\text{cone}_{\text{TC}}(X)$ .  $\square$

**Lemma 3.32.** *Suppose that  $V, W \subset \mathbb{T}_{\pm}^d$  are two nonempty finite sets. Let  $\hat{V} = \{(v, 0) \mid v \in V\}$ ,  $\hat{W} = \{(w, \mathbf{0}) \mid w \in W\}$ , and*

$$X = \text{conv}_{\text{TC}}(\{v \triangleleft \lambda \odot w \mid v \in V, w \in W, \lambda \in \mathbb{T}_{\geq \mathbf{0}}\}) .$$

*Then, we have the equality*

$$\{(x, 0) \mid x \in X\} = \text{cone}_{\text{TC}}(\hat{V} \cup \hat{W}) \cap \{x_{d+1} = 0\} .$$

*Proof.* Let  $x \in X$ . By Proposition 3.27, we have

$$x \in \text{Faces}(\mu_1 \odot (v_1 \triangleleft \lambda_1 \odot w_1), \dots, \mu_\ell \odot (v_\ell \triangleleft \lambda_\ell \odot w_\ell))$$

for some  $\mu_i \in \mathbb{T}_{\geq \mathbf{0}}$ ,  $\bigoplus_i \mu_i = \mathbf{0}$ ,  $\lambda_i \in \mathbb{T}_{\geq \mathbf{0}}$ ,  $v_i \in V$ ,  $w_i \in W$ . Denote  $\xi_i = \mu_i \odot \lambda_i$  for all  $i$  and observe that

$$\begin{aligned} & \text{Vert}(\mu_1 \odot (v_1 \triangleleft \lambda_1 \odot w_1), \dots, \mu_\ell \odot (v_\ell \triangleleft \lambda_\ell \odot w_\ell)) \\ &= \text{Vert}(\mu_1 \odot v_1 \triangleleft \xi_1 \odot w_1, \dots, \mu_\ell \odot v_\ell \triangleleft \xi_\ell \odot w_\ell) \\ &\subseteq \text{Vert}(\mu_1 \odot v_1, \dots, \mu_\ell \odot v_\ell, \xi_1 \odot w_1, \dots, \xi_\ell \odot w_\ell) . \end{aligned}$$

By Lemma 3.23 we get  $z \in \text{Faces}(\mu_1 \odot v_1, \dots, \mu_\ell \odot v_\ell, \xi_1 \odot w_1, \dots, \xi_\ell \odot w_\ell)$ . Let  $\hat{v}_i = (v_i, 0) \in \hat{V}$  and  $\hat{w}_i = (w_i, \mathbf{0}) \in \hat{W}$  for all  $i$ . Since  $\bigoplus_i \mu_i = \mathbf{0}$  we get

$$(x, 0) \in \text{Faces}(\mu_1 \odot \hat{v}_1, \dots, \mu_\ell \odot \hat{v}_\ell, \xi_1 \odot \hat{w}_1, \dots, \xi_\ell \odot \hat{w}_\ell) .$$

In particular, Lemma 3.30 implies that  $(x, 0) \in \text{cone}_{\text{TC}}(\hat{V} \cup \hat{W})$ . Conversely, suppose that  $(x, 0) \in \text{cone}_{\text{TC}}(\hat{V} \cup \hat{W})$ . Then, Lemma 3.30 shows that there exist  $\mu \in \mathbb{T}_{\geq \mathbf{0}}^\ell$ ,  $\xi \in \mathbb{T}_{\geq \mathbf{0}}^\ell$ ,  $\hat{v}_i \in \hat{V}$ ,  $\hat{w}_i \in \hat{W}$  such that  $(x, 0) \in \text{Faces}(\mu_1 \odot \hat{v}_1, \dots, \mu_\ell \odot \hat{v}_\ell, \xi_1 \odot \hat{w}_1, \dots, \xi_\ell \odot \hat{w}_\ell)$ . Since the last coordinate of  $(x, 0)$  is 0, we get  $\bigoplus_i \mu_i = \mathbf{0}$ . Without loss of generality, we can suppose that  $\mu_1 = \mathbf{0}$ . Observe that

$$\begin{aligned} & \text{Vert}(\mu_1 \odot \hat{v}_1, \dots, \mu_\ell \odot \hat{v}_\ell, \xi_1 \odot \hat{w}_1, \dots, \xi_\ell \odot \hat{w}_\ell) \\ &= \text{Vert}(\mu_1 \odot \hat{v}_1, \dots, \mu_\ell \odot \hat{v}_\ell, (-1) \odot \hat{v}_1 \triangleleft \xi_1 \odot \hat{w}_1, \dots, (-1) \odot \hat{v}_1 \triangleleft \xi_\ell \odot \hat{w}_\ell) \\ &= \text{Vert}(\mu_1 \odot \hat{v}_1, \dots, \mu_\ell \odot \hat{v}_\ell, (-1) \odot \hat{z}_1, \dots, (-1) \odot \hat{z}_\ell) , \end{aligned}$$

where  $\hat{z} = \hat{v}_1 \triangleleft (\xi_i + 1) \odot \hat{w}_i$ . Hence, by Lemma 3.23 we get

$$(x, 0) \in \text{Faces}(\mu_1 \odot \hat{v}_1, \dots, \mu_\ell \odot \hat{v}_\ell, (-1) \odot \hat{z}_1, \dots, (-1) \odot \hat{z}_\ell).$$

For every  $i$ , let  $v_i, z_i \in \mathbb{T}_\pm^d$  be the projection of  $\hat{v}_i, \hat{z}_i$  obtained by deleting the last coordinate. Then, we have

$$\{v_1, \dots, v_\ell, z_1, \dots, z_\ell\} \subseteq \{v \triangleleft \lambda \odot w \mid v \in V, w \in W, \lambda \in \mathbb{T}_{\geq 0}\}.$$

Furthermore, since the last coordinate of every point  $\hat{v}_i, \hat{z}_i$  is equal to 0 and  $\bigoplus_i \mu_i = 0$ , we get  $x \in \text{Faces}(\mu_1 \odot v_1, \dots, \mu_\ell \odot v_\ell, (-1) \odot z_1, \dots, (-1) \odot z_\ell)$ . Therefore, by Proposition 3.27 we have  $x \in X$ .  $\square$

#### 4. LIFTS OF SIGNED HALFSAPACES

Some of our results rely on a correspondence between tropical halfspaces and halfspaces defined over non-Archimedean valued fields. In the following, we consider the field of (*generalized*) *real Puiseux series*  $\mathbb{K} = \mathbb{R}\{t\}$ , whose elements

$$\gamma = \sum c_i t^{a_i}, \quad a_i, c_i \in \mathbb{R}, \quad a_0 > a_1 > a_2 > \dots \quad (23)$$

are formal power series with real exponents and such that the sequence  $(a_i)_i$  is either finite or unbounded. The addition and multiplication of Puiseux series are defined in the natural way. Furthermore, given a series  $\gamma$  as in (23), we say that  $c_0$  is its *leading coefficient* and we denote by  $\text{lc}: \mathbb{K} \rightarrow \mathbb{R}$  the map that sends a Puiseux series to its leading coefficient, with the convention that  $\text{lc}(0) = 0$ . We also say  $\gamma$  is *positive* if its leading coefficient is positive. This makes  $\mathbb{K}$  an ordered field via  $\gamma > \delta$  if and only if  $\gamma - \delta$  is positive. It is known that  $\mathbb{K}$  is a real closed field [45] and this remains true even if one considers a subfield formed by Puiseux series that are absolutely convergent for sufficiently large  $t$  [52]. All our results are valid for both of these fields. The crucial property of a real closed field is that for ‘well-structured’ statements (in the sense of model theory), they behave exactly as the ‘usual’ real numbers via Tarski’s principle [50], [44, Corollary 3.3.16].

The field of Puiseux series is linked with signed tropical numbers via the signed valuation map.

**Definition 4.1.** The map  $\text{sval}: \mathbb{K} \rightarrow \mathbb{T}_\pm$  sends a Puiseux series as in (23) to its *signed valuation*,

$$\text{sval}(\gamma) = \begin{cases} a_0 & \text{if } \gamma > 0 \\ \mathbb{0} & \text{if } \gamma = 0 \\ \ominus a_0 & \text{if } \gamma < 0 \end{cases}.$$

We extend  $\text{sval}$  to vectors in  $\mathbb{K}^d$  componentwise by putting  $\text{sval}(\mathbf{x}) = (\text{sval}(\mathbf{x}_i)_{i \in [d]})$ .

General properties of the images of semialgebraic sets under the signed valuation map were studied in [36], see also [13, Section 4].

The following lemma summarizes basic properties of the signed valuation.

**Lemma 4.2.** *The signed valuation map has the following properties:*

- (i) if  $\mathbf{x}_1, \mathbf{x}_2 \in \mathbb{K}$  satisfy  $\mathbf{x}_1 \geq \mathbf{x}_2$ , then  $\text{sval}(\mathbf{x}_1) \geq \text{sval}(\mathbf{x}_2)$ ;
- (ii) if  $\mathbf{x}_1, \dots, \mathbf{x}_n \in \mathbb{K}$ , then  $\text{sval}(\mathbf{x}_1 \cdots \mathbf{x}_n) = \text{sval}(\mathbf{x}_1) \odot \cdots \odot \text{sval}(\mathbf{x}_n)$ ;
- (iii) if  $\mathbf{x}_1, \dots, \mathbf{x}_n \in \mathbb{K}^d$ , then  $\text{sval}(\mathbf{x}_1 + \cdots + \mathbf{x}_n) \in \mathcal{U}(\text{sval}(\mathbf{x}_1) \oplus \cdots \oplus \text{sval}(\mathbf{x}_n))$ .

*Proof.* The first property is trivial if  $\mathbf{x}_1 \geq 0 \geq \mathbf{x}_2$ . Suppose that  $\mathbf{x}_1 \geq \mathbf{x}_2 > 0$  but  $\text{sval}(\mathbf{x}_1) < \text{sval}(\mathbf{x}_2)$ . Then, the leading coefficient of  $\mathbf{x}_2 - \mathbf{x}_1$  is equal to  $\text{lc}(\mathbf{x}_2) > 0$ , so  $\mathbf{x}_2 - \mathbf{x}_1 > 0$ , which gives a contradiction. Analogously, if  $0 > \mathbf{x}_1 \geq \mathbf{x}_2$ , but  $\text{sval}(\mathbf{x}_1) < \text{sval}(\mathbf{x}_2)$ , then the leading coefficient of  $\mathbf{x}_2 - \mathbf{x}_1$  is equal to  $-\text{lc}(\mathbf{x}_1) > 0$ , which gives a contradiction.

The second property for  $n = 2$  follows from the definition of multiplication of Puiseux series. The extension to  $n > 2$  follows by an immediate induction.

To prove the third property, it is enough to consider the case  $d = 1$  since  $\text{sval}$  and  $\mathcal{U}$  are defined componentwise. Furthermore, observe that if  $\mathbf{y}, \mathbf{z} \in \mathbb{K}$  are nonnegative (or nonpositive), then  $\text{sval}(\mathbf{y} + \mathbf{z}) = \text{sval}(\mathbf{y}) \oplus \text{sval}(\mathbf{z})$  by the definition of addition in Puiseux series. We can order  $\mathbf{x}_1, \dots, \mathbf{x}_n$  in such a way that  $\mathbf{x}_1, \dots, \mathbf{x}_k$  are nonnegative and  $\mathbf{x}_{k+1}, \dots, \mathbf{x}_n$  are negative. Let  $\mathbf{x}^+ = \mathbf{x}_1 + \dots + \mathbf{x}_k$  and  $\mathbf{x}^- = \mathbf{x}_{k+1} + \dots + \mathbf{x}_n$ . By the observation above, we have  $\text{sval}(\mathbf{x}^+) = \text{sval}(\mathbf{x}_1) \oplus \dots \oplus \text{sval}(\mathbf{x}_k)$  and  $\text{sval}(\mathbf{x}^-) = \text{sval}(\mathbf{x}_{k+1}) \oplus \dots \oplus \text{sval}(\mathbf{x}_n)$ . If  $\text{sval}(\mathbf{x}^+) \neq \ominus \text{sval}(\mathbf{x}^-)$ , then the definition of addition in Puiseux series implies that  $\text{sval}(\mathbf{x}^+ + \mathbf{x}^-) = \text{sval}(\mathbf{x}^+) \oplus \text{sval}(\mathbf{x}^-)$ . If  $\text{sval}(\mathbf{x}^+) = \ominus \text{sval}(\mathbf{x}^-) = a$ , then the definition of addition in Puiseux series implies that  $\text{sval}(\mathbf{x}^+ + \mathbf{x}^-) \in [\ominus a, a] = \mathcal{U}(\text{sval}(\mathbf{x}^+) \oplus \text{sval}(\mathbf{x}^-))$ .  $\square$

The signed valuation map allows us to study sets defined over  $\mathbb{K}^d$  by looking at their images under  $\text{sval}$ . Conversely, it is sometimes useful to study a set defined in  $\mathbb{T}_{\pm}^d$  by looking at its ‘lift’ in the Puiseux series. Since  $\text{sval}$  is not bijective, we have many possible choices for the lift. We now introduce different types of lifts of points  $x \in \mathbb{T}_{\pm}^d$  into Puiseux series that are used in this work. Similar lifts were used in [9] to derive properties about lifts of tropical halfspaces within  $\mathbb{T}_{\geq 0}^d$ . We start with the simplest one, the canonical lift.

**Definition 4.3.** Given a point  $x \in \mathbb{T}_{\pm}^d$  we define its *canonical lift*,  $\text{cli}(x) \in \mathbb{K}^d$ , as the point

$$\forall i \in [d], \quad (\text{cli}(x))_i = \sigma t^{|x_i|},$$

where  $\sigma \in \{-, 0, +\}$  is the ‘de-tropicalized’ version of  $\text{tsgn}(x_i) \in \{\ominus, \mathbf{0}, \oplus\}$  and we use the convention that  $t^{\mathbf{0}} = 0$ .

The canonical lifts are very simple, but it turns out that they are not particularly well suited for our purposes. Instead, it is more useful for us to consider lifts that vary from one orthant of  $\mathbb{T}_{\pm}^d$  to another. To this end, we introduce the following definition.

**Definition 4.4.** Given a point  $x \in \mathbb{T}_{\pm}^d$  and a set  $J \subseteq [d]$  we define the *lift of  $x$  of type  $J$* , denoted  $\text{li}_J(x) \in \mathbb{K}^d$ , by

$$\forall i \in [d], \quad (\text{li}_J(x))_i = \begin{cases} (d+1)t^{|x_i|} & \text{if } \text{tsgn}(x_i) = \oplus \text{ and } i \in J, \\ -(d+1)t^{|x_i|} & \text{if } \text{tsgn}(x_i) = \ominus \text{ and } i \notin J, \\ \sigma t^{|x_i|} & \text{otherwise,} \end{cases}$$

where  $\sigma$  is the ‘untropical’ version of  $\text{tsgn}(x_i)$ .

In this way,  $\text{li}_{\emptyset}(x)$  coincides with  $\text{cli}(x)$  on the nonnegative orthant of  $\mathbb{T}_{\pm}^d$ ,  $\text{li}_{[d]}(x)$  coincides with  $\text{cli}(x)$  on the nonpositive orthant of  $\mathbb{T}_{\pm}^d$  and so on. Also, for every  $x$  and  $J$  we have  $\text{sval}(\text{li}_J(x)) = x$ .

**4.1. Tropicalization of halfspaces.** We now characterize the signed valuation of halfspaces. This is a simple generalization of the characterization known for tropical halfspaces in one orthant. To start, we fix our notation for halfspaces over Puiseux series.

**Definition 4.5.** For a vector  $(\mathbf{a}_0, \mathbf{a}_1, \dots, \mathbf{a}_d) \in \mathbb{K}^{d+1}$  such that  $(\mathbf{a}_1, \dots, \mathbf{a}_d) \neq 0$  we define the (closed affine) halfspace by

$$\overline{\mathcal{H}}^+(\mathbf{a}) = \left\{ \mathbf{x} \in \mathbb{K}^d \mid \mathbf{a} \cdot \begin{pmatrix} 0 \\ \mathbf{x} \end{pmatrix} \geq 0 \right\}. \quad (24)$$

Furthermore, we denote  $\overline{\mathcal{H}}^-(\mathbf{a}) = \overline{\mathcal{H}}^+(-\mathbf{a})$ .

The next lemma characterizes the signed valuation of halfspaces.

**Lemma 4.6.** For every halfspace  $\overline{\mathcal{H}}^+(\mathbf{a})$  we have  $\text{sval}(\overline{\mathcal{H}}^+(\mathbf{a})) = \overline{\mathcal{H}}^+(\text{sval}(\mathbf{a}))$ .

*Proof.* The proof proceeds as in the case of tropical halfspaces in one orthant, cf. [29, Proposition 2.4]. Let  $a = \text{sval}(\mathbf{a})$ . The inclusion  $\text{sval}(\overline{\mathcal{H}}^+(\mathbf{a})) \subseteq \overline{\mathcal{H}}^+(a)$  follows from the arithmetic properties of  $\text{sval}$ . Indeed, if  $\mathbf{x} \in \mathbb{K}^d$  is such that  $\mathbf{y} = \mathbf{a}_0 + \mathbf{a}_1 \mathbf{x}_1 + \dots + \mathbf{a}_d \mathbf{x}_d \geq 0$  and we denote  $x = \text{sval}(\mathbf{x})$ , then Lemma 4.2 shows that  $\text{sval}(\mathbf{y}) \geq 0$  and  $\text{sval}(\mathbf{y}) \in \mathcal{U}(a_0 \oplus a_1 \odot x_1 \oplus \dots \oplus a_d \odot x_d)$ . Hence, we either have  $a_0 \oplus a_1 \odot x_1 \oplus \dots \oplus a_d \odot x_d \in \mathbb{T}_\bullet$  or  $a_0 \oplus a_1 \odot x_1 \oplus \dots \oplus a_d \odot x_d \in \mathbb{T}_{>0}$ .

Conversely, if  $x \in \mathcal{H}^+(a)$  and we let  $\mathbf{x} = \text{cli}(x)$ , then Lemma 4.2 shows that  $\text{sval}(\mathbf{a}_0 + \mathbf{a}_1 \mathbf{x}_1 + \dots + \mathbf{a}_d \mathbf{x}_d) \in \mathcal{U}(a_0 \oplus a_1 \odot x_1 \oplus \dots \oplus a_d \odot x_d)$ , which is a singleton in  $\mathbb{T}_{>0}$ , so  $\mathbf{x} \in \overline{\mathcal{H}}^+(\mathbf{a})$ . Hence  $\mathcal{H}^+(a) \subseteq \text{sval}(\overline{\mathcal{H}}^+(\mathbf{a})) \subseteq \overline{\mathcal{H}}^+(a)$ . Furthermore, the set  $\text{sval}(\overline{\mathcal{H}}^+(\mathbf{a}))$  is closed by [36, Theorem 6.9] or, equivalently, [13, Corollary 4.11]. Therefore, Lemma 2.4 implies that  $\text{sval}(\overline{\mathcal{H}}^+(\mathbf{a})) = \overline{\mathcal{H}}^+(a)$ .  $\square$

The next lemma gives a more explicit connection between tropical halfspaces of type  $J$  and lifts of type  $J$ .

**Lemma 4.7.** Suppose that  $\overline{\mathcal{H}}^+(a)$  is a tropical halfspace of type  $J \subseteq [d]$ . Let  $x \in \overline{\mathcal{H}}^+(a)$  and denote  $K = [d]_0 \setminus J$ . Then  $\text{li}_J(x) \in \overline{\mathcal{H}}^+(\text{li}_K(a))$ .

Before giving the proof, let us note that if  $x \in \mathcal{H}^+(a)$ , then any lift of  $x$  belongs to  $\overline{\mathcal{H}}^+(\mathbf{a})$  for any lift  $\mathbf{a}$  of  $a$ . However, in order to lift the points that belong to the boundary of  $\overline{\mathcal{H}}^+(a)$  we need to be more careful.

*Proof.* Let  $\mathbf{x} = \text{li}_J(x)$  and  $\mathbf{a} = \text{li}_K(a)$ . By definition, we have

$$\mathbf{a}_0 = \begin{cases} (d+1)t^{a_0} & \text{if } a_0 \in \mathbb{T}_{>0}, \\ -t^{|a_0|} & \text{otherwise.} \end{cases}$$

Also, for every  $i \in [d]$  we have  $\mathbf{a}_i = \text{tsgn}(a_i)t^{|a_i|}$  and

$$\mathbf{a}_i \mathbf{x}_i = \begin{cases} (d+1)t^{|a_i|+|x_i|} & \text{if } \text{tsgn}(x_i) = \text{tsgn}(a_i), \\ -t^{|a_i|+|x_i|} & \text{otherwise.} \end{cases}$$

Therefore, we have  $\mathbf{a} \cdot \begin{pmatrix} 0 \\ \mathbf{x} \end{pmatrix} \geq 0$  if and only if

$$(d+1) \left( \alpha t^{|a_0|} + \sum_{\text{tsgn}(x_i) = \text{tsgn}(a_i)} t^{|a_i|+|x_i|} \right) \geq \beta t^{|a_0|} + \sum_{\text{tsgn}(x_i) \neq \text{tsgn}(a_i)} t^{|a_i|+|x_i|}, \quad (25)$$



where  $(\alpha, \beta) = (1, 0)$  if  $a_0 \in \mathbb{T}_{>0}$  and  $(\alpha, \beta) = (0, 1)$  otherwise. Let  $\mathbf{y} \in \mathbb{K}$  denote the series on the left-hand side of (10) and  $\mathbf{z} \in \mathbb{K}$  denote the series on the right-hand side of (10). By Lemma 4.2 we have  $\text{sval}(\mathbf{y} - \mathbf{z}) \in \mathcal{U}(\text{sval}(\mathbf{y}) \ominus \text{sval}(\mathbf{z})) = \mathcal{U}(a_0 \oplus a_1 \odot x_1 \oplus \cdots \oplus a_d \odot x_d)$ . If  $\mathcal{U}(a_0 \oplus a_1 \odot x_1 \oplus \cdots \oplus a_d \odot x_d)$  is a nonnegative singleton, then  $\mathbf{y} - \mathbf{z} \geq 0$ . Otherwise, we have  $\text{sval}(\mathbf{y}) = \text{sval}(\mathbf{z}) > 0$ . In this case, note that  $\text{lc}(\mathbf{y}) \geq d + 1$  and  $\text{lc}(\mathbf{z}) \leq d$ . Hence  $\mathbf{y} - \mathbf{z} \geq 0$ .  $\square$

**4.2. Separation over Puiseux series.** We also need the following version of the hyperplane separation theorem over Puiseux series. We recall that a set  $\mathbf{X} \subseteq \mathbb{K}^d$  is a *cone* if  $\lambda \mathbf{x} \in \mathbf{X}$  for all  $\mathbf{x} \in \mathbf{X}$  and  $\lambda > 0$ . We also recall that a set  $\mathbf{X}$  is *semialgebraic* if it is defined by a finite Boolean combination of polynomial inequalities. Since the field of Puiseux series is real closed, the semialgebraic sets over  $\mathbb{K}^d$  have similar properties to the semialgebraic sets over  $\mathbb{R}^d$ —we refer to [18] for more information on this topic. For the sake of generality, we state the next two propositions for semialgebraic sets, but the familiarity with semialgebraic sets is not necessary to understand the other results of this paper—it is enough to admit that polyhedra are semialgebraic. The first proposition is a hyperplane separation theorem for convex semialgebraic sets.

**Proposition 4.8.** *Suppose that  $\mathbf{X}, \mathbf{Y} \subseteq \mathbb{K}^d$  are two nonempty convex semialgebraic sets such that  $\mathbf{X} \cap \mathbf{Y} = \emptyset$ . Then, there exists a halfspace  $\overline{\mathcal{H}}^+(\mathbf{a})$  such that  $\mathbf{X} \subseteq \overline{\mathcal{H}}^+(\mathbf{a})$  and  $\mathbf{Y} \subseteq \overline{\mathcal{H}}^-(\mathbf{a})$ .*

*Proof.* If we replace  $\mathbb{K}$  by  $\mathbb{R}$ , then the claim follows from the hyperplane separation theorem in  $\mathbb{R}^d$ , see, e.g., [47, Theorem 11.3 and Theorem 11.7]. Since the sets  $\mathbf{X}, \mathbf{Y}$  are supposed to be semialgebraic, the claim for  $\mathbb{K}$  follows from the completeness of the theory of real closed fields, see, e.g., [44, Corollary 3.3.16] or [18, Theorem 2.80]. The argument is based on the existence of finite formulas for describing  $\mathbf{X}$  and  $\mathbf{Y}$  and the existence of a separating hyperplane in a real closed field due to their convexity; for more details see Appendix A.  $\square$

*Remark 4.9.* We note that the assumption that  $\mathbf{X}, \mathbf{Y}$  are semialgebraic cannot be entirely skipped. Indeed, it is shown in [46] that  $\mathbb{R}$  is the only ordered field that admits the general hyperplane separation theorem. For the interested reader, we adapt the example from [46] to  $\mathbb{K}$  in Appendix A. We also note that a special case of Proposition 4.8 is still valid for sets that are definable in definably complete extensions of real closed fields, see [16, Corollary 2.20]. Also, [46] gives a version of the separation theorem that is valid for arbitrary ordered fields.

The next proposition is an application of Proposition 4.8 that characterizes the signed valuations.

**Theorem 4.10.** *Suppose that  $\mathbf{X} \subseteq \mathbb{K}^d$  is a nonempty closed convex semialgebraic set. Then, for every  $y \notin \text{sval}(\mathbf{X})$  there exists  $\mathbf{a} \in \mathbb{K}^{d+1}$  such that  $\mathbf{X} \subseteq \overline{\mathcal{H}}^+(\mathbf{a})$  and  $y \notin \overline{\mathcal{H}}^+(\text{sval}(\mathbf{a}))$ . In particular,  $\text{sval}(\mathbf{X})$  is equal to the intersection of the closed tropical halfspaces that contain it.*

*Proof.* To prove the first part of the claim, let  $y \notin \text{sval}(\mathbf{X})$ . Since  $\mathbf{X}$  is closed and semialgebraic,  $\text{sval}(\mathbf{X})$  is also closed by [36, Theorem 6.9] or [13, Corollary 4.11]. Therefore, there exists an open neighborhood of  $y$  that does not intersect  $\text{sval}(\mathbf{X})$ . Since the order on  $\mathbb{T}_\pm$  is dense, we can find  $\ell_1, \dots, \ell_d, r_1, \dots, r_d \in \mathbb{T}_\pm$  such that

$$\ell_1 < y_1 < r_1, \ell_2 < y_2 < r_2, \dots, \ell_d < y_d < r_d$$

and such that the box  $B = [\ell_1, r_1] \times [\ell_2, r_2] \times \cdots \times [\ell_d, r_d]$  does not intersect  $\text{sval}(\mathbf{X})$ . Consider the lifted box

$$\mathbf{B} = [\text{cli}(\ell_1), \text{cli}(r_1)] \times [\text{cli}(\ell_2), \text{cli}(r_2)] \times \cdots \times [\text{cli}(\ell_d), \text{cli}(r_d)] \subset \mathbb{K}^d.$$

The set  $\mathbf{B}$  is convex and semialgebraic. Moreover, we have  $\mathbf{X} \cap \mathbf{B} = \emptyset$  because  $\text{sval}(\mathbf{X}) \cap \text{sval}(\mathbf{B}) = \text{sval}(\mathbf{X}) \cap B = \emptyset$ . Hence, by Proposition 4.8, there exists  $\mathbf{a} \in \mathbb{K}^{d+1}$  such that  $\mathbf{X} \subseteq \overline{\mathcal{H}}^+(\mathbf{a})$  and  $\mathbf{B} \subseteq \overline{\mathcal{H}}^-(\mathbf{a})$ . Therefore  $B \subseteq \overline{\mathcal{H}}^-(\text{sval}(\mathbf{a}))$  by Lemma 4.6. Since  $y$  belongs to the interior of  $B$ , Lemma 2.4 shows that  $y \in \mathcal{H}^-(\text{sval}(\mathbf{a}))$ . In particular, we have  $y \notin \overline{\mathcal{H}}^+(\text{sval}(\mathbf{a}))$ . To prove the second part of the claim, note that  $\text{sval}(\mathbf{X}) \subseteq \overline{\mathcal{H}}^+(\text{sval}(\mathbf{a}))$  by Lemma 4.6. In other words, the closed tropical halfspace  $\overline{\mathcal{H}}^+(\text{sval}(\mathbf{a}))$  contains  $\text{sval}(\mathbf{X})$  but not  $y$ . Since  $y$  was arbitrary, we get that  $\text{sval}(\mathbf{X})$  is an intersection of some family of closed tropical halfspaces. Therefore, it is also an intersection of all the closed tropical halfspaces that contain it.  $\square$

**Corollary 4.11.** *If  $\mathbf{X} \subseteq \mathbb{K}^d$  from Theorem 4.10 is a cone, we can choose the tropical halfspaces to be linear.*

*Proof.* To see that, suppose that  $\mathbf{a}_0 \neq 0$ . Since  $0 \in \mathbf{X}$ , we have  $\mathbf{a}_0 > 0$ . Consider  $\hat{\mathbf{a}} = (0, \mathbf{a}_1, \dots, \mathbf{a}_d)$ . Since  $\overline{\mathcal{H}}^+(\hat{\mathbf{a}}) \subseteq \overline{\mathcal{H}}^+(\mathbf{a})$ , we get  $y \notin \overline{\mathcal{H}}^+(\text{sval}(\hat{\mathbf{a}}))$  by Lemma 4.6.

It remains to show that  $\mathbf{X} \subseteq \overline{\mathcal{H}}^+(\hat{\mathbf{a}})$ . Indeed, if there exists  $\mathbf{x} \in \mathbf{X}$  such that  $\mathbf{x}^T \hat{\mathbf{a}} < 0$ , then for any  $\lambda > -\mathbf{a}_0/\mathbf{x}^T \hat{\mathbf{a}} > 0$  we have  $\lambda \mathbf{x} \in \mathbf{P}$  but  $\lambda \mathbf{x} \notin \overline{\mathcal{H}}^+(\mathbf{a})$ , which is a contradiction.

Therefore, we can suppose that  $\mathbf{a}$  satisfies  $\mathbf{a}_0 = 0$ .  $\square$

*Example 4.12.* We note that neither of the assumptions of Theorem 4.10 can be skipped. Indeed, the set  $\mathbf{X} = \{\mathbf{x} \in \mathbb{K} \mid \mathbf{x} > 0\}$  is semialgebraic and convex but not closed. We have  $\text{sval}(\mathbf{X}) = \mathbb{T}_{>0}$ , which is not an intersection of closed tropical halfspaces. Likewise, the set  $\mathbf{Y} = \{\mathbf{x} \in \mathbb{K} \mid \text{sval}(\mathbf{x}) > 0\}$  is closed and convex, but not semialgebraic. We have  $\text{sval}(\mathbf{Y}) = \{x \in \mathbb{T}_{\pm} \mid x > 0\}$ , which is not an intersection of closed tropical halfspaces.

The following lemma gives a partial characterization of the intersection of all closed tropical halfspaces that contain a given finite set.

**Lemma 4.13.** *Given a finite set  $X = \{x_1, \dots, x_m\} \subset \mathbb{T}_{\pm}^d$  we have*

$$\begin{aligned} \bigcap \left\{ \overline{\mathcal{H}}^+(a) \mid X \subseteq \overline{\mathcal{H}}^+(a) \right\} &= \bigcap \left\{ \text{sval}(\mathbf{X}) \mid \mathbf{X} \subseteq \mathbb{K}^d \text{ convex} \wedge X \subseteq \text{sval}(\mathbf{X}) \right\} \\ &= \bigcap_{J \subseteq [d]} \text{sval} \left( \text{conv}(\text{li}_J(x_1), \dots, \text{li}_J(x_m)) \right). \end{aligned}$$

*Proof.* Denote

$$\begin{aligned} U &= \bigcap \left\{ \overline{\mathcal{H}}^+(a) \mid X \subseteq \overline{\mathcal{H}}^+(a) \right\}, \\ V &= \bigcap \left\{ \text{sval}(\mathbf{X}) \mid \mathbf{X} \subseteq \mathbb{K}^d \text{ convex} \wedge X \subseteq \text{sval}(\mathbf{X}) \right\}, \\ W &= \bigcap_{J \subseteq [d]} \text{sval} \left( \text{conv}(\text{li}_J(x_1), \dots, \text{li}_J(x_m)) \right). \end{aligned}$$

We start by showing that

$$V = \bigcap \left\{ \text{sval}(\text{conv}(\mathbf{x}_1, \dots, \mathbf{x}_m)) \mid \forall i, \text{sval}(\mathbf{x}_i) = x_i \right\}. \quad (26)$$

Indeed, the inclusion  $\subseteq$  in (26) holds as we just range over a smaller set. To prove the opposite inclusion, it is enough to observe that any convex set  $\mathbf{X}$  such that  $\{x_1, \dots, x_m\} \subseteq \text{sval}(\mathbf{X})$  contains some set of the form  $\text{conv}(\mathbf{x}_1, \dots, \mathbf{x}_m)$ . Hence, the equality (26) holds.

As  $\text{conv}(\mathbf{x}_1, \dots, \mathbf{x}_m)$  is a closed convex semialgebraic set for each choice with  $\text{sval}(\mathbf{x}_i) = x_i$  for all  $i \in [m]$ , we can apply Theorem 4.10 to each set in the intersection (26). This implies that  $V$  is an intersection of a family of closed tropical halfspaces. Since  $X \subseteq V$ , all of these halfspaces contain  $X$ . Hence,  $U \subseteq V$ .

The inclusion  $V \subseteq W$  is trivial. Therefore, it remains to prove that  $W \subseteq U$ . To this end, take a point  $z \notin U$ . By definition, there is a closed tropical halfspace  $\overline{\mathcal{H}}^+(a)$  that contains  $X$  with  $z \notin \overline{\mathcal{H}}^+(a)$ . Suppose that  $\overline{\mathcal{H}}^+(a)$  is of type  $J \subseteq [d]$  and let  $K = [d]_0 \setminus J$  be its complement. Then, Lemma 4.7 shows that  $\text{li}_J(x_i) \in \overline{\mathcal{H}}^+(\text{li}_K(a))$  for every  $i \in [m]$ . Since the set  $\overline{\mathcal{H}}^+(\text{li}_K(a))$  is convex, we obtain

$$\text{conv}(\text{li}_J(x_1), \dots, \text{li}_J(x_m)) \subset \overline{\mathcal{H}}^+(\text{li}_K(a)).$$

Using the representation of the valuation of halfspace from Lemma 4.6, we get

$$W \subseteq \text{sval}(\text{conv}(\text{li}_J(x_1), \dots, \text{li}_J(x_m))) \subset \text{sval}(\overline{\mathcal{H}}^+(\text{li}_K(a))) = \overline{\mathcal{H}}^+(a).$$

In particular, we get  $z \notin W$  and so  $W \subseteq U$  as claimed.  $\square$

The next lemma strengthens the claim of Corollary 4.11 for polyhedral cones.

**Lemma 4.14.** *Suppose that  $\mathbf{P} \subseteq \mathbb{K}^d$  is a polyhedral cone. Then,  $\text{sval}(\mathbf{P})$  is an intersection of finitely many linear closed tropical halfspaces.*

*Proof.* Let  $\mathbf{P} = \{\mathbf{x} \in \mathbb{K}^d \mid \mathbf{A}\mathbf{x} \geq 0\}$  for some matrix  $\mathbf{A} \in \mathbb{K}^{n \times d}$ . By Farkas' lemma [49, Corollary 7.1h], for every  $\mathbf{a} \in \mathbb{K}^d$  we have the equivalence

$$\mathbf{P} \subseteq \overline{\mathcal{H}}^+(0, \mathbf{a}) \iff \mathbf{a} \in \{\mathbf{y}^T \mathbf{A} \mid \mathbf{y} \geq 0\}. \quad (27)$$

The intersection of  $\{\mathbf{y}^T \mathbf{A} \mid \mathbf{y} \geq 0\}$  with any closed orthant  $\mathbb{K}^d$  is a polyhedral cone, and so it is generated by a finite family of rays by the Minkowski–Weyl theorem [49, Corollary 7.1a]. For any closed orthant  $\mathbf{O} \subset \mathbb{K}^d$ , let  $\mathbf{U}_{\mathbf{O}} \subset \mathbb{K}^d$  be a finite set such that  $\{\mathbf{y}^T \mathbf{A} \mid \mathbf{y} \geq 0\} \cap \mathbf{O} = \text{cone}(\mathbf{U}_{\mathbf{O}})$ . Let  $\mathbf{U} = \bigcup_{\mathbf{O}} \mathbf{U}_{\mathbf{O}}$  be the set of all rays obtained in this way. We claim that  $\text{sval}(\mathbf{P}) = \bigcap_{\mathbf{u} \in \mathbf{U}} \overline{\mathcal{H}}^+(\mathbf{O}, \text{sval}(\mathbf{u}))$ . The inclusion “ $\subseteq$ ” follows by combining (27) with Lemma 4.6.

To prove the opposite inclusion, let  $y \notin \text{sval}(\mathbf{P})$ . By Theorem 4.10, there exists  $\mathbf{a} \in \mathbb{K}^{d+1}$  such that  $\mathbf{P} \subseteq \overline{\mathcal{H}}^+(\mathbf{a})$  and  $y \notin \overline{\mathcal{H}}^+(\text{sval}(\mathbf{a}))$ . Since  $\mathbf{P}$  is a cone, we can choose  $\mathbf{a}$  in such a way that  $\mathbf{a}_0 = 0$  by Corollary 4.11; hence, we let  $\mathbf{a} \in \mathbb{K}^d$ .

Let  $\mathbf{O} \subset \mathbb{K}^d$  be a closed orthant such that  $\mathbf{a} = (\mathbf{a}_1, \dots, \mathbf{a}_d) \in \mathbf{O}$ . By (27) we have  $\mathbf{a} \in \text{cone}(\mathbf{U}_{\mathbf{O}})$ . Denote  $\mathbf{U}_{\mathbf{O}} = \{\mathbf{u}_1, \dots, \mathbf{u}_m\}$ . Since the orthant  $\mathbf{O}$  is fixed, we have  $\text{sval}(\mathbf{a}) \in \text{cone}_{\text{TC}}(\text{sval}(\mathbf{U}_{\mathbf{O}}))$  by [8, Lemma 8]. Suppose that  $y \in \overline{\mathcal{H}}^+(\mathbf{O}, \text{sval}(\mathbf{u}_i))$  for all  $i \in [m]$ . Then, we also have  $\text{sval}(\mathbf{u}_i) \in \overline{\mathcal{H}}^+(\mathbf{O}, y)$  for all  $i \in [m]$ . As closed halfspaces are convex by Corollary 3.8, we get  $\text{conv}_{\text{TC}}(\text{sval}(\mathbf{U}_{\mathbf{O}})) =$

$\text{conv}_{\text{TC}}(\text{sval}(\mathbf{u}_1), \dots, \text{sval}(\mathbf{u}_m)) \subseteq \overline{\mathcal{H}}^+(\mathbb{O}, y)$ . Furthermore, since the tropical halfspace  $\overline{\mathcal{H}}^+(\mathbb{O}, y)$  is linear, we get  $\text{conv}_{\text{TC}}(\text{sval}(\mathbf{U}\mathbf{O})) \subseteq \overline{\mathcal{H}}^+(\mathbb{O}, y)$ . Therefore  $\text{sval}(\mathbf{a}) \in \overline{\mathcal{H}}^+(\mathbb{O}, y)$ , which implies that  $y \in \overline{\mathcal{H}}^+(\mathbb{O}, \text{sval}(\mathbf{a}))$ , giving a contradiction.

Hence, there exists  $i \in [m]$  such that  $y \notin \overline{\mathcal{H}}^+(\mathbb{O}, \text{sval}(\mathbf{u}_i))$ . Since  $y$  was arbitrary, we get  $\bigcap_{\mathbf{u} \in \mathbf{U}} \overline{\mathcal{H}}^+(\mathbb{O}, \text{sval}(\mathbf{u})) \subseteq \text{sval}(\mathbf{P})$ .  $\square$

**Lemma 4.15.** *Let  $\mathbf{X} \subseteq \mathbb{K}^d$  be an arbitrary set. Take  $\lambda \in \mathbb{K}$  such that  $\lambda \geq 0$  and denote  $\text{sval}(\lambda) = \lambda$ . Then, we have*

$$\text{sval}(\text{conv}(\lambda\mathbf{X})) = \lambda \odot \text{sval}(\text{conv}(\mathbf{X})). \quad (28)$$

In particular, we have the equality

$$\text{sval}(\text{cone}(\mathbf{X})) = \bigcup \{ \text{sval}(\text{conv}(t^\lambda \mathbf{X})) \mid \lambda \in \mathbb{T}_{\geq \mathbb{O}} \}.$$

*Proof.* Let  $x \in \text{sval}(\text{conv}(\lambda\mathbf{X}))$ . Since  $\text{conv}(\lambda\mathbf{X}) = \lambda \text{conv}(\mathbf{X})$ , there exists  $\mathbf{y} \in \text{conv}(\mathbf{X})$  such that  $x = \text{sval}(\lambda\mathbf{y}) = \text{sval}(\lambda) \odot \text{sval}(\mathbf{y}) = \lambda \odot \text{sval}(\mathbf{y})$ . Hence  $x \in \lambda \odot \text{sval}(\text{conv}(\mathbf{X}))$ . Conversely, suppose that  $x \in \lambda \odot \text{sval}(\text{conv}(\mathbf{X}))$ . Then, there exists  $\mathbf{y} \in \text{conv}(\mathbf{X})$  such that  $x = \lambda \odot \text{sval}(\mathbf{y}) = \text{sval}(\lambda\mathbf{y})$ , so that  $x \in \text{sval}(\text{conv}(\lambda\mathbf{X}))$ . To prove the second claim, note that (28) gives  $\text{sval}(\text{conv}(\lambda\mathbf{X})) = \text{sval}(\text{conv}(t^\lambda \mathbf{X}))$ . In particular, we get

$$\begin{aligned} \text{sval}(\text{cone}(\mathbf{X})) &= \bigcup \{ \text{sval}(\text{conv}(\lambda\mathbf{X})) \mid \lambda \geq 0 \} \\ &= \bigcup \{ \text{sval}(\text{conv}(t^\lambda \mathbf{X})) \mid \lambda \in \mathbb{T}_{\geq \mathbb{O}} \}. \end{aligned} \quad \square$$

## 5. SEPARATION AND HEMISPACES

A *TC-hemisphere* is a TC-convex subset  $X \subseteq \mathbb{T}_{\pm}^d$  for which also  $\mathbb{T}_{\pm}^d \setminus X$  is TC-convex. The fundamental elimination property of TC-convexity (Theorem 5.2) leads to a representation of a TC-convex set as intersection of its containing TC-hemispheres (Theorem 5.3). Then, we identify that TC-hemispheres are close to halfspaces (Theorem 5.10); based on a more thorough study of TC-hemispheres, this leads to a representation of TC-hemispheres by convex lifts (Proposition 5.18).

**5.1. Fundamental separation property.** For  $\sim \in \{\leq, \geq\}^d$ , a point  $w$  with  $y_k \sim_k w_k$  for all  $k \in [d]$  is said to *dominate*  $y$  with respect to  $\sim$ .

**Lemma 5.1.** *If  $\mu \leq 0$  and  $w \in \mathcal{U}(v^1 \oplus \mu \odot v^2)$ , then  $\text{Faces}(X \cup v^1) \cap \text{Faces}(Y \cup v^2) \subseteq \text{Faces}(X \cup \mu \odot Y \cup w)$ .*

*Proof.* Let  $z$  be an arbitrary point in  $\text{Faces}(X \cup v^1) \cap \text{Faces}(Y \cup v^2)$ . Using Corollary 3.21, for every  $\sim \in \{\leq, \geq\}^d$ , there exist vertices  $\tilde{x}_1 \triangleleft v^1 \triangleleft \tilde{x}_2$  and  $\tilde{y}_1 \triangleleft v^2 \triangleleft \tilde{y}_2$  in  $\text{Vert}(X \cup v^1)$  and  $\text{Vert}(Y \cup v^2)$ , respectively, which dominate  $z$  with respect to  $\sim$ . Here,  $\tilde{x}_1, \tilde{x}_2$  arise as left sum of points in  $X$  and  $\tilde{y}_1, \tilde{y}_2$  arise analogously. As  $\tilde{x}_1 \triangleleft \mu \odot \tilde{y}_1 \in \mathcal{U}(\tilde{x}_1 \oplus \mu \odot \tilde{y}_1)$ ,  $w \in \mathcal{U}(v^1 \oplus \mu \odot v^2)$  and  $\tilde{x}_2 \triangleleft \mu \odot \tilde{y}_2 \in \mathcal{U}(\tilde{x}_2 \oplus \mu \odot \tilde{y}_2)$ , by Lemma 2.15,  $\tilde{x}_1 \triangleleft \mu \odot \tilde{y}_1 \triangleleft w \triangleleft \tilde{x}_2 \triangleleft \mu \odot \tilde{y}_2$  dominates  $z$  with respect to  $\sim$ . Ranging over all  $\sim$ , these points form a subset of  $\text{Vert}(X \cup \mu \odot Y \cup w)$ . Again using Corollary 3.21,  $z$  is contained in  $\text{Faces}(X \cup \mu \odot Y \cup w)$ .  $\square$

**Theorem 5.2.** *Fix a set  $X = \{x_0, \dots, x_n\} \subset \mathbb{T}_{\pm}^d$ . Then, for every  $y_1, y_2, z \in \mathbb{T}_{\pm}^d$  such that  $x_0 \in \text{conv}_{\text{TO}}(y_1, y_2)$  we have the implication*

$$(z \in \text{conv}_{\text{TC}}(X \cup \{y_1\}) \wedge z \in \text{conv}_{\text{TC}}(X \cup \{y_2\})) \implies z \in \text{conv}_{\text{TC}}(X).$$

*Proof.* Breaking the symmetry between  $y_1$  and  $y_2$ , there is a  $\mu \in \mathbb{T}_{\geq 0}$  with  $\mu \leq 0$  such that  $x_0 \in \mathcal{U}(y_1 \oplus \mu \odot y_2)$ . So let  $\lambda^{(1)} \in \mathbb{T}_{\geq 0}^{n+1}$ ,  $\lambda^{(2)} \in \mathbb{T}_{\geq 0}^{n+1}$  and  $\rho_1, \rho_2 \in \mathbb{T}_{\geq 0}$  with  $\bigoplus \lambda^{(1)} \oplus \rho_1 = \bigoplus \lambda^{(2)} \oplus \rho_2 = 0$ .

$$z \in \text{Faces}(\text{diag}(\lambda^{(1)}) \odot X \cup \rho_1 \odot y_1) \cap \text{Faces}(\text{diag}(\lambda^{(2)}) \odot X \cup \rho_2 \odot y_2).$$

First, if  $\rho_2 = \mathbf{0}$  then  $z \in \text{Faces}(\text{diag}(\lambda^{(2)}) \odot X)$  yields the desired containment. Furthermore, if  $\mu = \mathbf{0}$  then  $x_0 = y_1$  which concludes the claim with  $z \in \text{Faces}(\text{diag}(\lambda^{(1)}) \odot X \cup \rho_1 \odot y_1)$ .

Otherwise, let  $\delta = \rho_1 - \rho_2 + \mu = \rho_1 \odot \rho_2^{\odot -1} \odot \mu$ , or equivalently  $\delta \odot \rho_2 = \rho_1 \odot \mu$ . We have to distinguish two cases, depending on the real sign of  $\delta$ .

**Case 1** ( $\delta \leq 0$ .) By Lemma 5.1, we obtain that

$$z \in \text{Faces}(\text{diag}(\lambda^{(1)}) \odot X \cup \delta \odot \text{diag}(\lambda^{(2)}) \odot X \cup w)$$

for an arbitrary  $w \in \mathcal{U}(\rho_1 \odot y_1 \oplus \delta \odot \rho_2 \odot y_2)$ . With  $\delta \odot \rho_2 = \rho_1 \odot \mu$ , this implies that

$$z \in \text{Faces}(\text{diag}(\lambda^{(1)}) \odot X \cup \delta \odot \text{diag}(\lambda^{(2)}) \odot X \cup \rho_1 \odot x_0).$$

In particular, this yields  $z \in \text{conv}_{\text{TC}}(X)$  since  $\delta, \rho_1 \leq 0$  and, therefore,  $\bigoplus \lambda^{(1)} \oplus \delta \odot \bigoplus \lambda^{(2)} \oplus \rho_1 = \bigoplus \lambda^{(1)} \oplus \rho_1 = 0$ .

**Case 2** ( $\delta > 0$ .) By Lemma 5.1, we obtain that

$$z \in \text{Faces}(\delta^{\odot -1} \odot \text{diag}(\lambda^{(1)}) \odot X \cup \text{diag}(\lambda^{(2)}) \odot X \cup w)$$

for an arbitrary  $w \in \mathcal{U}(\delta^{\odot -1} \odot \rho_1 \odot y_1 \oplus \rho_2 \odot y_2)$ . With  $\delta^{\odot -1} \odot \rho_1 = \rho_2 \odot \mu^{\odot -1}$ , this implies that

$$z \in \text{Faces}(\delta^{\odot -1} \odot \text{diag}(\lambda^{(1)}) \odot X \cup \text{diag}(\lambda^{(2)}) \odot X \cup \rho_2 \odot \mu^{\odot -1} \odot x_0).$$

In particular, this yields  $z \in \text{conv}_{\text{TC}}(X)$  since  $\delta^{\odot -1}, \rho_2 \leq \rho_2 \odot \mu^{\odot -1} = \delta^{\odot -1} \odot \rho_1 < 0$  and, therefore,

$$\bigoplus \lambda^{(2)} \leq \delta^{\odot -1} \odot (\bigoplus \lambda^{(1)}) \oplus \bigoplus \lambda^{(2)} \oplus \rho_2 \odot \mu^{\odot -1} \leq \bigoplus \lambda^{(2)} \oplus \rho_2 = 0 = \bigoplus \lambda^{(2)},$$

using  $\delta^{\odot -1} \odot (\bigoplus \lambda^{(1)}) < 0$  and  $\rho_2 \odot \mu^{\odot -1} < 0$ .  $\square$

The maximal set of elements which is tropically convex and does not contain a fixed point is called *semispace* in [39]. This concept (for unsigned tropical convexity) was used to study hemispaces. We use a similar idea in the following proof.

**Theorem 5.3.** *Every TC-convex set  $G \subseteq \mathbb{T}_{\pm}^d$  is an intersection of TC-hemispaces.*

*More precisely, each of these TC-hemispaces can be chosen such that its complement is TO-convex.*

*Proof.* The proof is a variant of the argument given in [40, Thm. 5.2] and [51, Thm. I.4.13]. If  $G \in \{\emptyset, \mathbb{T}_{\pm}^d\}$ , then the claim is trivial. Otherwise, let  $z \notin G$ . We want to prove that there exists a TC-hemisphere  $H^*$  such that  $G \subseteq H^*$  and  $z \notin H^*$ . To do so, we consider the family  $\mathcal{F}$  of TC-convex subsets  $S$  of  $\mathbb{T}_{\pm}^d$  such that  $G \subseteq S$  and  $z \notin S$ . We partially order  $\mathcal{F}$  by inclusion. Any chain  $\mathcal{C} \subseteq \mathcal{F}$  is upper bounded by  $(\bigcup_{S \in \mathcal{C}} S) \in \mathcal{F}$  using that TC-convexity is closed under arbitrary nested unions. Therefore, the Kuratowski–Zorn lemma implies that  $\mathcal{F}$  has at least one maximal element  $H^*$ . By construction,  $H^*$  is TC-convex; hence to prove that  $H^*$  is a TC-hemisphere we show that  $\mathbb{T}_{\pm}^d \setminus H^*$  is TO-convex. Note that this implies that it is also TC-convex. Suppose that  $\mathbb{T}_{\pm}^d \setminus H^*$  is not TO-convex. Then, there exist  $y_1, y_2 \in \mathbb{T}_{\pm}^d \setminus H^*$  and  $x_0 \in \text{conv}_{\text{TO}}(y_1, y_2)$  such that  $x_0 \in H^*$ . Furthermore, by

the maximality of  $H^*$  we get  $z \in \text{conv}_{\text{TC}}(H^* \cup \{y_1\})$  and  $z \in \text{conv}_{\text{TC}}(H^* \cup \{y_2\})$ . Hence, by Proposition 3.24, there exist finite sets  $X_1, X_2 \subseteq H^*$  such that  $z \in \text{conv}_{\text{TC}}(X_1 \cup \{y_1\})$  and  $z \in \text{conv}_{\text{TC}}(X_2 \cup \{y_2\})$ . By putting  $X = \{x_0\} \cup X_1 \cup X_2$ , we get  $x_0 \in \text{conv}_{\text{TO}}(y_1, y_2)$ ,  $z \in \text{conv}_{\text{TC}}(X \cup \{y_1\})$ ,  $z \in \text{conv}_{\text{TC}}(X \cup \{y_2\})$ , and  $z \notin \text{conv}_{\text{TC}}(X) \subseteq H^*$ . This gives a contradiction with Theorem 5.2. Since  $z \notin G$  was arbitrary, we obtain that  $G$  is an intersection of TC-hemispace.  $\square$

**5.2. TC-hemispace are nearly halfspaces.** For a cone  $X \subseteq \mathbb{T}_{\pm}^d$ , a *relative conic TC-hemisphere w.r.t.  $X$*  is a subset  $H \subseteq X \subseteq \mathbb{T}_{\pm}^d$  such that  $H$  and  $X \setminus H$  are TC-convex cones.

Note that if  $G \subseteq \mathbb{T}_{\pm}^d$  is a TC-hemisphere as well as a cone and  $X \subseteq \mathbb{T}_{\pm}^d$  is a TC-convex cone then  $G \cap X$  is a relative conic TC-hemisphere w.r.t.  $X$ .

We summarize a crucial insight on the structure of tropical hemispace in the non-negative orthant  $\mathbb{T}_{\geq 0}^d$ ; see [22], [39, Section 4], [30, Section 4].

**Proposition 5.4.** *If  $G \subseteq \mathbb{T}_{> 0}^d$  is a relative conic TC-hemisphere w.r.t.  $\mathbb{T}_{> 0}^d$  and  $G \notin \{\emptyset, \mathbb{T}_{> 0}^d\}$ , then there exists a vector  $(a_1, \dots, a_d) \in \mathbb{T}_{\pm}^d$ ,  $(a_1, \dots, a_d) \neq \mathbf{0}$ , such that  $\mathcal{H}^+(\mathbf{0}, a) \cap \mathbb{T}_{> 0}^d \subseteq G \subseteq \overline{\mathcal{H}^+(\mathbf{0}, a)} \cap \mathbb{T}_{> 0}^d$ .*

Proposition 5.4 can be extended to characterize relative hemispace that are included in the nonnegative orthant and contain (parts of) the boundary of  $\mathbb{T}_{\geq 0}^d$ . The next corollary gives one such extension that is needed in our proofs.

**Corollary 5.5.** *Let  $X = \{x \in \mathbb{T}_{\geq 0}^d \mid x_1 > \mathbf{0}\}$ . If  $G \subseteq X$  is a relative conic TC-hemisphere w.r.t.  $X$  and  $G \cap \mathbb{T}_{> 0}^d \notin \{\emptyset, \mathbb{T}_{> 0}^d\}$ , then there exists a vector  $(a_1, \dots, a_d) \in \mathbb{T}_{\pm}^d$ ,  $(a_1, \dots, a_d) \neq \mathbf{0}$ , such that  $\mathcal{H}^+(\mathbf{0}, a) \cap X \subseteq G \subseteq \overline{\mathcal{H}^+(\mathbf{0}, a)} \cap X$ .*

*Proof.* By Proposition 5.4, there exists a vector  $(a_1, \dots, a_d) \in \mathbb{T}_{\pm}^d$ ,  $(a_1, \dots, a_d) \neq \mathbf{0}$ , such that  $\mathcal{H}^+(\mathbf{0}, a) \cap \mathbb{T}_{> 0}^d \subseteq G \cap \mathbb{T}_{> 0}^d \subseteq \overline{\mathcal{H}^+(\mathbf{0}, a)} \cap \mathbb{T}_{> 0}^d$ . Let  $x \in \mathcal{H}^+(\mathbf{0}, a) \cap X$  be any point and let  $y \in \mathbb{T}_{> 0}^d$  be such that  $y \notin G$ . Then, for sufficiently small  $\omega \in \mathbb{T}_{> 0}$ , the point  $x \triangleleft \omega \odot y \in \mathbb{T}_{> 0}^d$  belongs to  $\mathcal{H}^+(\mathbf{0}, a)$ , which implies that it belongs to  $G$ . Hence, by the TC-convexity of  $X \setminus G$ , we have  $x \in G$ . This shows that  $\mathcal{H}^+(\mathbf{0}, a) \cap X \subseteq G$ . Analogously, we get  $\mathcal{H}^-(\mathbf{0}, a) \cap X \subseteq (X \setminus G)$ , which implies that  $G \subseteq \overline{\mathcal{H}^+(\mathbf{0}, a)} \cap X$ .  $\square$

We note that the assumption  $G \cap \mathbb{T}_{> 0}^d \notin \{\emptyset, \mathbb{T}_{> 0}^d\}$  implies that  $\text{supp}^{\oplus}(a) \neq \emptyset$  and  $\text{supp}^{\ominus}(a) \neq \emptyset$ .

In order to prove our characterization of TC-hemispace we need to generalize Corollary 5.5 to handle multiple orthants. We start by characterizing relative hemispace w.r.t. the halfspace  $X = \{x \in \mathbb{T}_{\pm}^d \mid x_1 > \mathbf{0}\}$ . To do so, in Lemmas 5.6 to 5.8 we suppose that  $G$  is a relative conic TC-hemisphere w.r.t.  $X$ . Furthermore, we suppose that there exists an orthant  $O$  in  $\text{cl}(X)$  such that  $G \cap \text{int}(O) \notin \{\emptyset, \text{int}(O)\}$ . Then,  $G \cap O$  is a relative conic TC-hemisphere w.r.t.  $O \cap X$ . In particular, by Corollary 5.5, there is a vector  $(a_1, \dots, a_d) \in \mathbb{T}_{\pm}^d$  with  $(a_1, \dots, a_d) \neq \mathbf{0}$  such that  $\mathcal{H}^+(\mathbf{0}, a) \cap O \cap X \subseteq G \cap O \subseteq \overline{\mathcal{H}^+(\mathbf{0}, a)} \cap O \cap X$ . Let  $Q$  be a neighboring orthant of  $O$  in  $X$ , i.e., an orthant obtained from  $O$  by changing one sign (other than the sign of the first coordinate). In the next two lemmas (Lemma 5.6 and Lemma 5.7), we show that the relative TC-hemispace  $G \cap O$  in  $O$  and  $G \cap Q$  in  $Q$  are essentially determined by the same vector  $a$ .

By suitably flipping signs and permuting variables, we can assume that  $O = \mathbb{T}_{\geq 0}^d$  and  $Q$  differs in the sign of the  $d$ -th component. We denote  $T = (O \cup Q) \cap X$ .

In the following proofs,  $\omega \in \mathbb{T}_{> 0}$  and  $\Omega \in \mathbb{T}_{> 0}$  will mean a sufficiently small and sufficiently big number, respectively.

**Lemma 5.6.** *There are exactly two possibilities for the part of the relative TC-hemisphere  $G$  in  $Q$  to be trivial:*

- (i)  $G \cap \text{int}(Q) = \text{int}(Q) \Leftrightarrow \text{supp}^\ominus(a) = \{d\} \Leftrightarrow \{y \in Q \cap X \mid y_d < \mathbb{O}\} \subseteq G$ ,
- (ii)  $G \cap \text{int}(Q) = \emptyset \Leftrightarrow \text{supp}^\oplus(a) = \{d\} \Leftrightarrow \{y \in Q \cap X \mid y_d < \mathbb{O}\} \cap G = \emptyset$ .

*Proof.* To start, suppose that  $G \cap \text{int}(Q) = \text{int}(Q)$  but there is  $k \in \text{supp}^\ominus(a) \cap [d-1]$ . As we assume that  $G \cap \text{int}(O) \not\subseteq \{\emptyset, \text{int}(O)\}$ , there exists  $j \in \text{supp}^\oplus(a)$ . We define  $x \in O, y \in Q$  and derive  $x \triangleleft y \in O$  via

$$x_i = \begin{cases} \Omega & i = j \\ \omega & \text{else} \end{cases} \quad y_i = \begin{cases} 2\Omega & i = k \\ \ominus\omega & i = d \\ \omega & \text{else} \end{cases} \quad (x \triangleleft y)_i = \begin{cases} 2\Omega & i = k \\ \Omega & i = j \\ \omega & \text{else} \end{cases}$$

By construction,  $x, y \in G$  but  $x \triangleleft y \notin G$ , as one can see from  $a \odot x > \mathbb{O}$  and  $a \odot (x \triangleleft y) < \mathbb{O}$ . This contradicts the fact that  $G$  is TC-convex. Hence,  $\text{supp}^\ominus(a) \subseteq \{d\}$  and as  $G \cap \text{int}(O) \not\subseteq \{\emptyset, \text{int}(O)\}$  we get  $\text{supp}^\ominus(a) = \{d\}$ .

Now, assume that  $\text{supp}^\ominus(a) = \{d\}$  but there is a point  $y \in (Q \cap X) \setminus G$  such that  $y_d < \mathbb{O}$ . Consider two cases: if  $y_k > \mathbb{O}$  for some  $k \in \text{supp}^\oplus(a)$ , then we define  $x, z \in O \cap X$  via

$$x_i = \begin{cases} \Omega & i = d \\ y_i & \text{else} \end{cases} \quad z_i = \begin{cases} \omega & i = d \\ y_i & \text{else} \end{cases}$$

Then  $z$  is on the line segment between  $x$  and  $y$ , but  $x, y \in T \setminus G$  while  $z \in G$ . This contradicts the fact that  $G$  is a relative TC-hemisphere. If  $y_k = \mathbb{O}$  for every  $k \in \text{supp}^\oplus(a)$ , then we define  $x, z \in O \cap X$  via

$$x_i = \begin{cases} |y_d| & i = d \\ \omega & \text{else} \end{cases} \quad z_i = \begin{cases} 2\omega & i = d \\ y_i & i \neq d, y_i \neq \mathbb{O} \\ \omega & \text{else} \end{cases}$$

We note that  $z \in \text{Faces}(x, y)$ . Moreover, the assumption on  $y$  implies that  $a \odot z > \mathbb{O}$ . Hence, as above we have  $x, y \in T \setminus G$  while  $z \in G$ , contradicting the fact that  $G$  is a relative TC-hemisphere. The implication  $\{y \in Q \cap X \mid y_d < \mathbb{O}\} \subseteq G \Rightarrow G \cap \text{int}(Q) = \text{int}(Q)$  is trivial, proving the first point of the lemma.

The second point follows analogously as it can be obtained by considering the complement of  $G$ .  $\square$

**Lemma 5.7.**  $\mathcal{H}^+(\mathbb{O}, a) \cap T \subseteq G \cap T \subseteq \overline{\mathcal{H}^+}(\mathbb{O}, a) \cap T$ .

*Proof.* The case of a trivial intersection of  $G$  with  $\text{int}(Q)$  follows directly from Lemma 5.6. Hence, by Corollary 5.5, we can assume that there exists a vector  $(b_1, \dots, b_d) \in \mathbb{T}_{\pm}^d \setminus \{\mathbb{O}\}$  such that  $\mathcal{H}^+(\mathbb{O}, b) \cap Q \cap X \subseteq G \cap Q \subseteq \overline{\mathcal{H}^+}(\mathbb{O}, b) \cap Q \cap X$ . We show with an exhaustive case distinction that the defining vectors  $a$  and  $b$  agree up to positive scaling; then this concludes the proof of the lemma.

**Case 1** (Different sign pattern.)

Let  $j$  be the smallest index for which  $\text{tsgn}(a_j) \neq \text{tsgn}(b_j)$ . Breaking the symmetry in  $a$  and  $b$ , we can assume that  $j \in \text{supp}(a)$ .

By taking the complement in both orthants, we can assume that  $j \in \text{supp}^\oplus(a)$  which means  $\text{tsgn}(b_j) \in \{\mathbb{O}, \ominus\}$ .

**Case 1a** ( $j = d$ .)

Since the sign of  $a$  and  $b$  differ in  $d = j \in \text{supp}^\oplus(a)$ , there is an  $r \in \text{supp}^\oplus(b) \setminus \{d\}$ . As the relative TC-hemisphere in  $\text{int}(O)$  is not trivial, there is an index  $s \in \text{supp}^\ominus(a)$ . Note that  $r, s, d$  are pairwise different due to the minimality assumption on  $j$ . We define  $x \in O, y \in Q$  and derive  $x \oplus y$  via

$$x_i = \begin{cases} \Omega & i = s \\ 2\Omega & i = d \\ \omega & \text{else} \end{cases} \quad y_i = \begin{cases} 0 & i = r \\ \ominus 2\Omega & i = d \\ \omega & \text{else} \end{cases} \quad (x \oplus y)_i = \begin{cases} 0 & i = r \\ \Omega & i = s \\ \bullet 2\Omega & i = d \\ \omega & \text{else} \end{cases}.$$

By construction,  $a \odot x > \mathbb{O}, b \odot y > \mathbb{O}$ . Choosing  $z \in \mathcal{U}(x \oplus y)$  with  $z_d = \omega$ , we get a point in  $G$ . However,  $z \in O$  and  $a \odot z < \mathbb{O}$ , a contradiction.

**Case 1b** ( $j \neq d$ .)

Let  $k = \min \text{supp}^\ominus(a)$  (which cannot equal  $j$  then). As the relative TC-hemisphere in  $\text{int}(Q)$  is not trivial, there is an index  $r \in \text{supp}^\ominus(b)$  (which could equal  $j$ ).

We define  $x \in O, y \in Q$  and derive  $x \triangleleft y \in O$  via

$$x_i = \begin{cases} 0 & i = k \\ \omega & \text{else} \end{cases} \quad y_i = \begin{cases} \Omega & i = j \\ 0 & i = r, i \neq j \\ \ominus \omega & i = d \\ \omega & \text{else} \end{cases} \quad (x \triangleleft y)_i = \begin{cases} \Omega & i = j \\ 0 & i = r, i \neq j \\ 0 & i = k \\ \omega & \text{else} \end{cases}$$

By construction,  $a \odot x < \mathbb{O}, b \odot y < \mathbb{O}$  but  $a \odot (x \triangleleft y) > \mathbb{O}$ , a contradiction to the TC-convexity of  $G$ .

From the cases considered so far, we deduce that  $a$  and  $b$  have the same sign pattern. Hence, only the following possibility is remaining.

**Case 2** Now, we have that the sign patterns of  $a$  and  $b$  are the same.

If  $a$  and  $b$  are not the same up to scaling, then there are three indices  $i, j, k \in [d]$  such that not all three signs of the respective components agree and among the quotients  $a_i \odot b_i^{\ominus-1}, a_j \odot b_j^{\ominus-1}, a_k \odot b_k^{\ominus-1}$  at least one has a different value from the other two. Then there are indices  $p, q \in \{i, j, k\}$  with  $\text{tsgn}(a_p) = \text{tsgn}(b_p) = \oplus$ ,  $\text{tsgn}(a_q) = \text{tsgn}(b_q) = \ominus$ , and  $a_p \odot b_q \neq a_q \odot b_p$ .

**Case 2a** ( $d \notin \{p, q\}$ .)

By scaling  $a$  and  $b$ , we can assume that  $a_p = b_p = 0$ . By the symmetry in  $a$  and  $b$ , we can assume that  $|a_q| > |b_q|$ .

We choose  $\xi_a, \xi_b \in \mathbb{T}_{>\mathbb{O}}$  with  $\xi_a < |a_q^{\ominus-1}| < \xi_b < |b_q^{\ominus-1}|$ . With this, we define  $x \in O, y \in Q$  and derive  $x \triangleleft y \in O$  via

$$x_i = \begin{cases} \xi_a & i = q \\ 0 & i = p \\ \omega & \text{else} \end{cases} \quad y_i = \begin{cases} \xi_b & i = q \\ \ominus \omega & i = d \\ 0 & i = p \\ \omega & \text{else} \end{cases} \quad (x \triangleleft y)_i = \begin{cases} \xi_b & i = q \\ 0 & i = p \\ \omega & \text{else} \end{cases}$$

The choice of  $\xi_a$  and  $\xi_b$  yields  $|\xi_a \odot a_q| < 0$  and  $|\xi_b \odot b_q| < 0$  as well as  $|\xi_b \odot a_q| > 0$ . This implies that  $a \odot x > \mathbb{O}, b \odot y > \mathbb{O}$  and  $a \odot (x \triangleleft y) < \mathbb{O}$ , a contradiction.

**Case 2b** ( $d \in \{p, q\}$ .)



By taking complements in both orthants, we can assume that  $d = p$ , so  $d \in \text{supp}^\oplus(a) = \text{supp}^\oplus(b)$ . Since  $\overline{\mathcal{H}}^+(\mathbb{O}, b) \cap \text{int}(Q) \neq \emptyset$ , there is at least one  $r \in \text{supp}^\oplus(b) \setminus \{d\}$ . By scaling, we can assume that  $a_q = b_q = \ominus 0$ . By the previous case, we have  $a_r = b_r$ . By the symmetry in  $a$  and  $b$ , we can suppose that  $|a_d| > |b_d|$ . We choose  $\xi_r, \xi_q \in \mathbb{T}_{>0}$  with

$$a_d > \xi_q > \xi_r \odot a_r = \xi_r \odot b_r > b_d .$$

With this, we define  $x \in O, y \in Q$  and  $z \in O$  via

$$x_i = \begin{cases} \xi_q & i = q \\ 0 & i = d \\ \omega & \text{else} \end{cases} \quad y_i = \begin{cases} \xi_r & i = r \\ \ominus 0 & i = d \\ \omega & \text{else} \end{cases} \quad z_i = \begin{cases} \xi_q & i = q \\ \xi_r & i = r \\ \omega & \text{else} \end{cases}$$

We have  $z \in \text{Faces}(x, y)$ . Moreover,  $a \odot x > \mathbb{O}$ ,  $b \odot y > \mathbb{O}$ , but  $a \odot z < \mathbb{O}$ , a contradiction.  $\square$

We now extend Lemma 5.7 from a union of two orthants to the entire set  $X$ .

**Lemma 5.8.** *We have  $\mathcal{H}^+(\mathbb{O}, a) \cap X \subseteq G \subseteq \overline{\mathcal{H}}^+(\mathbb{O}, a) \cap X$ .*

*Proof.* Let  $Q$  be any orthant in  $\text{cl}(X)$ . We want to show that  $\mathcal{H}^+(a) \cap Q \cap X \subseteq G \cap Q \subseteq \overline{\mathcal{H}}^+(a) \cap Q \cap X$ . Let  $\nu \in \{\oplus, \ominus\}^d$  be the sign vector corresponding to  $Q$ , i.e., the vector such that  $\text{int}(Q) = \{x \in \mathbb{T}_\pm^d \mid \forall k, \text{tsgn}(x_k) = \nu_k\}$ . For the purposes of this proof, we say that an orthant is *good* if  $\mathcal{H}^+(\mathbb{O}, a)$  subdivides its interior in a non-trivial way; that is if there exists a pair  $(k, l) \in \text{supp}(a)$  such that  $\nu_k \text{tsgn}(a_k) \neq \nu_l \text{tsgn}(a_l)$ . Let  $r = |\{k \in [d] \mid k \in \text{supp}(a), \nu_k = \ominus\}|$ . We divide the proof into two cases.

**Case 1** (We have  $|\text{supp}(a)| \geq 3$  or  $r \in \{0, 1\}$ .)

In this case, we start by proving that there exists a sequence of orthants  $\mathbb{T}_{\geq 0}^d = Q_0, Q_1, \dots, Q_p = Q$  in  $\text{cl}(X)$  such that  $Q_i, Q_{i+1}$  differ by flipping one sign and the  $Q_i$  are good for all  $i \leq p-1$ . Such a sequence can be obtained in the following way. The orthant  $Q_0$  is good by the assumption of the lemma. To go from  $Q_0$  to  $Q$ , we first flip the signs in  $\{k \notin \text{supp}(a) \mid \nu_k = \ominus\}$  (in any order) and note that all orthants obtained in this way are good. Then, we flip the signs in  $\{k \in \text{supp}(a) \mid \nu_k = \ominus\}$ . If  $r \in \{0, 1\}$ , then this already proves the existence of the sequence, because this step either does nothing or flips one sign. If  $|\text{supp}(a)| \geq 3$  and  $r \geq 3$ , then there are at most two orthants that are not good and could be obtained at this step. However, since the graph of the  $r$ -dimensional hypercube is  $r$ -vertex-connected and  $r \geq 3$ , there is a way of flipping the signs in  $\{k \in \text{supp}(a) \mid \nu_k = \ominus\}$  that avoids going through these two orthants (except, possibly, for the last step, since  $Q$  may be not good). If  $|\text{supp}(a)| \geq 3$  and  $r = 2$ , then we let  $k, l$  be the two indices in  $\text{supp}(a)$  such that  $\nu_k = \nu_l = \ominus$  and  $j$  be any index in  $\text{supp}(a)$  such that  $\nu_j = \oplus$ . If  $\text{tsgn}(a_j) = \text{tsgn}(a_k)$  or  $\text{tsgn}(a_k) = \text{tsgn}(a_l)$ , then we first flip the sign of the component indexed by  $k$  and subsequently we flip the sign of the component indexed by  $l$ . If  $\text{tsgn}(a_j) = \text{tsgn}(a_l)$ , then we start by flipping the sign of the component indexed by  $l$  and subsequently we flip the sign of the component indexed by  $k$ .

Given the sequence  $Q_0, \dots, Q_p$  we apply Lemma 5.7 to  $T = (Q_0 \cup Q_1) \cap X$ . This gives  $\mathcal{H}^+(a) \cap Q_1 \cap X \subseteq G \cap Q_1 \subseteq \overline{\mathcal{H}}^+(a) \cap Q_1 \cap X$ . Furthermore, since  $Q_1$  is good,

Lemma 5.6 implies that  $G \cap \text{int}(Q_1) \neq \{\emptyset, \text{int}(Q_1)\}$ . Hence, by applying Lemmas 5.6 and 5.7 to  $T = (Q_1 \cup Q_2) \cap X$  we get that  $\mathcal{H}^+(a) \cap Q_2 \cap X \subseteq G \cap Q_2 \subseteq \overline{\mathcal{H}^+}(a) \cap Q_2 \cap X$  and  $G \cap \text{int}(Q_2) \neq \{\emptyset, \text{int}(Q_2)\}$ . By repeating this reasoning, we obtain the claim.

**Case 2** (We have  $|\text{supp}(a)| = r = 2$ .)

In this case, let  $\{k, l\} = \text{supp}(a)$  and let  $Q_{\{k\}}, Q_{\{l\}}, Q_{\{k,l\}}$  be the orthants obtained from  $\mathbb{T}_{\geq 0}^d$  by flipping the signs of the components indexed by  $k$  and  $l$ . Since  $\mathbb{T}_{\geq 0}^d$  is good, we can suppose (up to permuting  $k$  and  $l$ ) that  $\text{tsgn}(a_k) = \oplus$  and  $\text{tsgn}(a_l) = \ominus$ . Then, Lemma 5.6 shows that  $G \cap \text{int}(Q_{\{k\}}) = \emptyset$  and  $G \cap \text{int}(Q_{\{l\}}) = \text{int}(Q_{\{l\}})$ .

We first show that  $G \cap \text{int}(Q_{\{k,l\}}) \not\subseteq \{\emptyset, \text{int}(Q_{\{k,l\}})\}$ . We define  $x \in Q_{\{k,l\}}, y \in \mathbb{T}_{\geq 0}^d$  and derive  $x \triangleleft y \in Q_{\{k\}}$  via

$$x_i = \begin{cases} \ominus\omega & i = k \\ \ominus\Omega & i = l \\ 0 & \text{else} \end{cases} \quad y_i = \begin{cases} 0 & i = k \\ \Omega & i = l \\ 0 & \text{else} \end{cases} \quad (x \triangleleft y)_i = \begin{cases} 0 & i = k \\ \ominus\Omega & i = l \\ 0 & \text{else} \end{cases}$$

We have  $y \notin G$  and  $x \triangleleft y \in G$ . Therefore,  $x \in G \cap \text{int}(Q_{\{k,l\}})$ . Similarly, if we define

$$x_i = \begin{cases} \ominus\Omega & i = k \\ \ominus\omega & i = l \\ 0 & \text{else} \end{cases} \quad y_i = \begin{cases} \Omega & i = k \\ 0 & i = l \\ 0 & \text{else} \end{cases} \quad (x \triangleleft y)_i = \begin{cases} \ominus\Omega & i = k \\ 0 & i = l \\ 0 & \text{else} \end{cases}$$

then  $y \in G$  and  $x \triangleleft y \notin G$ . Hence, we have  $x \notin G$  and  $x \in \text{int}(Q_{\{k,l\}})$ . This implies  $G \cap \text{int}(Q_{\{k,l\}}) \not\subseteq \{\emptyset, \text{int}(Q_{\{k,l\}})\}$ .

Now, by Corollary 5.5, there exists a vector  $b \in \mathbb{T}_{\pm}^d \setminus \{\mathbf{0}\}$  such that  $\mathcal{H}^+(\mathbf{0}, b) \cap Q_{\{k,l\}} \subseteq G \cap Q_{\{k,l\}} \subseteq \overline{\mathcal{H}^+}(\mathbf{0}, b) \cap Q_{\{k,l\}}$ . By applying Lemma 5.6 to  $Q_{\{k,l\}} \cup Q_{\{k\}}$  and  $Q_{\{k,l\}} \cup Q_{\{l\}}$  we get  $\text{supp}^{\ominus}(b) = \text{supp}^{\ominus}(a) = \{l\}$  and  $\text{supp}^{\oplus}(b) = \text{supp}^{\oplus}(a) = \{k\}$ . By scaling  $a$  and  $b$ , we can assume that  $a_k = b_k = 0$ . We want to show that  $a_l = b_l$ . For contradiction, assume that  $|a_l| < \eta^{\ominus-1} < \xi^{\ominus-1} < |b_l|$  for some  $\eta, \xi \in \mathbb{T}_{>0}$ . We define  $x \in Q_{\{k,l\}}, y \in \mathbb{T}_{\geq 0}^d$  and derive  $x \triangleleft y \in Q_{\{k\}}$  via

$$x_i = \begin{cases} \ominus 0 & i = k \\ \ominus \xi & i = l \\ \omega & \text{else} \end{cases} \quad y_i = \begin{cases} 0 & i = k \\ \eta & i = l \\ \omega & \text{else} \end{cases} \quad (x \triangleleft y)_i = \begin{cases} \ominus 0 & i = k \\ \eta & i = l \\ \omega & \text{else} \end{cases} \quad (29)$$

We have  $x, y \in G$  (as  $b \odot x > \mathbf{0}$  and  $a \odot y > \mathbf{0}$ ) but  $x \triangleleft y \notin G$ , which gives a contradiction. Analogously, if  $|a_l| > \eta^{\ominus-1} > \xi^{\ominus-1} > |b_l|$  for some  $\xi, \eta \in \mathbb{T}_{>0}$ , and we define  $x, y$  as in (29), then  $x, y \notin G$  but  $y \triangleleft x \in G$ , giving a contradiction. Therefore, we have  $a = b$ .

To finish the proof, note that we can go from  $Q_{\{k,l\}}$  to  $Q$  by a sequence of good orthants obtained by flipping the signs that do not belong to  $\text{supp}(a)$ . This gives the claim by the same reasoning as in Case 1.  $\square$

*Example 5.9.* We illustrate the last case in the proof of Lemma 5.8.

Let  $a = (0, \ominus 0)$  and  $b = (0, \ominus 3)$ . We take  $\eta = -1$ ,  $\xi = -2$ , so  $x = (\ominus 0, \ominus(-2))$ ,  $y = (0, -1)$ . Then  $a \odot y = 0$ ,  $b \odot x = 1$  so  $x, y \in G$  but  $x \triangleleft y = (\ominus 0, -1)$  so  $a \odot (x \triangleleft y) = \ominus 0$ ,  $b \odot (x \triangleleft y) = \ominus 2$  and  $x \triangleleft y \notin G$ .




FIGURE 6. Sketch of the positions for the first part of Example 5.9.

Likewise, if  $a = (0, \ominus 3)$  and  $b = (0, \ominus 0)$ , then we take  $\eta = -2, \xi = -1$ , so  $x = (\ominus 0, \ominus(-1)), y = (0, -2)$ . We have  $a \odot y = \ominus 1, b \odot x = \ominus 0$  so  $x, y \notin G$  but  $y \triangleleft x = (0, \ominus(-1))$  satisfies  $a \odot (y \triangleleft x) = 2, b \odot (y \triangleleft x) = 0$  and  $y \triangleleft x \in G$ .

Now, we combine the last lemmas to prove that a TC-hemisphere is nearly a halfspace.

**Theorem 5.10.** *If  $G \subseteq \mathbb{T}_{\pm}^d$  is a TC-hemisphere and  $G \notin \{\emptyset, \mathbb{T}_{\pm}^d\}$ , then there exists a vector  $(a_0, \dots, a_d) \in \mathbb{T}_{\pm}^{d+1}$ ,  $(a_1, \dots, a_d) \neq \mathbf{0}$ , such that  $\mathcal{H}^+(a) \subseteq G \subseteq \overline{\mathcal{H}^+(a)}$ .*

*Proof.* We consider two cases.

**Case 1** (There exists an orthant  $O \in \mathbb{T}_{\pm}^d$  such that  $G \cap \text{int}(O) \notin \{\emptyset, \text{int}(O)\}$ .)

By flipping signs, we assume that  $O = \mathbb{T}_{>0}^d$ . Let  $X = \{x \in \mathbb{T}_{\pm}^{d+1} \mid x_{d+1} > \mathbf{0}\}$ . We define two sets  $H, \overline{H} \subseteq X$  by putting  $H = \text{cone}_{\text{TC}}\{(z, 0) \mid z \in G\} \setminus \{\mathbf{0}\}$  and  $\overline{H} = \text{cone}_{\text{TC}}\{(z, 0) \mid z \notin G\} \setminus \{\mathbf{0}\}$ . Since both  $G$  and its complement are TC-convex, we have

$$\begin{aligned} H &= \{\lambda \odot (z, 0) \mid \lambda \in \mathbb{T}_{>0}, z \in G\}, \\ \overline{H} &= \{\lambda \odot (z, 0) \mid \lambda \in \mathbb{T}_{>0}, z \notin G\}. \end{aligned}$$

In particular,  $H = \text{cone}_{\text{TC}}\{(z, 0) \mid z \in G\} \cap X$ , which implies that  $H$  is a TC-convex cone. Analogously,  $\overline{H}$  is a TC-convex cone.

Furthermore, we have  $H \cup \overline{H} = X$  because any point  $x = (x_1, \dots, x_{d+1})$  in  $X$  can be written as  $x = x_{d+1} \odot (y_1, \dots, y_d, 0)$ , where  $y_i = x_i \odot x_{d+1}^{\ominus 1}$  and  $y$  either belongs to  $G$  or not. Likewise, we have  $H \cap \overline{H} = \emptyset$  because if  $x \in H \cap \overline{H}$ , then the point  $y$  defined as above belongs both to  $G$  and its complement, which gives a contradiction. Thus,  $H$  is a relative conic TC-hemisphere w.r.t.  $X$  and  $H \cap \mathbb{T}_{>0}^{d+1} \notin \{\emptyset, \mathbb{T}_{>0}^{d+1}\}$ .

Hence, by Lemma 5.8, there exists a vector  $(a_1, \dots, a_{d+1}) \in \mathbb{T}_{\pm}^{d+1}$  such that  $(a_1, \dots, a_{d+1}) \neq \mathbf{0}$  and  $\mathcal{H}^+(a) \cap X \subseteq H \subseteq \overline{\mathcal{H}^+(a)} \cap X$ . In particular,

$$\begin{aligned} \{x \in \mathbb{T}_{\pm}^d \mid a_{d+1} \oplus a_1 \odot x_1 \oplus \dots \oplus a_d \odot x_d > \mathbf{0}\} &\subseteq G \\ &\subseteq \{x \in \mathbb{T}_{\pm}^d \mid a_{d+1} \oplus a_1 \odot x_1 \oplus \dots \oplus a_d \odot x_d \neq \mathbf{0}\}. \end{aligned}$$

Since  $\text{supp}^{\ominus}(a) \neq \emptyset$  and  $\text{supp}^{\oplus}(a) \neq \emptyset$ , we have  $(a_1, \dots, a_d) \neq \mathbf{0}$ .

**Case 2** ( $G \cap \text{int}(O) \in \{\emptyset, \text{int}(O)\}$  for every orthant  $O \subseteq \mathbb{T}_{\pm}^d$ .)

By taking suitable combinations, we see that  $G$  cannot only consist of parts of coordinate hyperplanes. Hence, by flipping signs, we can assume that  $\mathbb{T}_{>0}^d \subseteq G$ . We show that there is a  $k \in [d]$  such that

$$\{x \in \mathbb{T}_{\pm}^d \mid x_k > \mathbf{0}\} \subseteq G \subseteq \{x \in \mathbb{T}_{\pm}^d \mid x_k \geq \mathbf{0}\}. \quad (30)$$

Suppose the right inclusion does not hold, then for each  $\ell \in [d]$  there is an orthant  $O^{(\ell)}$  with  $\text{int}(O^{(\ell)}) \subseteq G$  whose  $\ell$ -th component is negative. As each  $O^{(\ell)}$  contains points close to the negative of the  $\ell$ -th tropical unit vector, taking convex combinations of these points and  $(0, \dots, 0) \in \mathbb{T}_{>0}^d$  yields points in the interior of every orthant in  $\mathbb{T}_{\pm}^d$ . This would imply  $G = \mathbb{T}_{\pm}^d$ , which was excluded. Hence,  $G \subseteq \{x \in \mathbb{T}_{\pm}^d \mid x_k \geq \mathbf{0}\}$ . Therefore, for half of the orthants the interior is contained in  $G$  and, by the same argument, for half of the orthants the interior is contained in the complement  $G'$  of  $G$ . But as  $G$  and  $G'$  cover the whole space, we get that  $G' \subseteq \{x \in \mathbb{T}_{\pm}^d \mid x_k \leq \mathbf{0}\}$ . By taking again the complement, this concludes the second case.  $\square$

As TO-hemispace are also TC-hemispace we immediately get the following.

**Corollary 5.11.** *If  $G \subseteq \mathbb{T}_{\pm}^d$  is a TO-hemispace and  $G \notin \{\emptyset, \mathbb{T}_{\pm}^d\}$ , then there exists a vector  $(a_0, \dots, a_d) \in \mathbb{T}_{\pm}^{d+1}$ ,  $(a_1, \dots, a_d) \neq \mathbf{0}$ , such that  $\mathcal{H}^+(a) \subseteq G \subseteq \overline{\mathcal{H}^+(a)}$ .*

**5.3. The boundary of TC-hemispace.** Before stating the next lemmas, we introduce the following notation. If  $a = (a_0, \dots, a_d) \in \mathbb{T}_{\pm}^{d+1}$  and  $x \in \mathbb{T}_{\pm}^d$ , then we denote

$$\begin{aligned} \text{Argmax}(a, x) &= \{k \in [d]_0 \mid a_k \odot x_k \neq \mathbf{0} \wedge \forall \ell \in [d]_0, |a_k| + |x_k| \geq |a_\ell| + |x_\ell|\} \\ &= \text{argmax}_{k \in [d]_0} |a_k \odot x_k| \cap \text{supp}(a) \cap \text{supp}(x) \end{aligned}$$

and

$$\text{Argmax}^+(a, x) = \{k \in \text{Argmax}(a, x) \mid a_k \odot x_k > \mathbf{0}\},$$

where we use the convention that  $x_0 = 0$ . Note that

$$\text{Argmax}^+(a, x) \subseteq \text{Argmax}(a, x) \subseteq \text{supp}(x) \cup \{0\}.$$


For a fixed  $a$ , comprising the points with (1) the same set  $\text{Argmax}(a, x)$ , (2) in the same orthant yields a cell decomposition of  $\mathbb{T}_{\pm}^d$ . Taking the common refinement of the cell decompositions for several possible  $a$  yields a generalization of the decomposition which was studied under the name ‘type decomposition’ [28] or ‘covector decomposition’ [38].

Right from the definition, we obtain some basic properties.

**Corollary 5.12.** *We have*

- (i) *If  $0 \notin \text{Argmax}(a, x) \cup \text{Argmax}(a, \rho \odot x)$ , then  $\text{Argmax}(a, \rho \odot x) = \text{Argmax}(a, x)$  for  $\rho \in \mathbb{T}_{\pm} \setminus \{\mathbf{0}\}$ ;*
- (ii)  *$\text{Argmax}(a, x) = \emptyset$  if and only if  $a \odot (0, x) = \mathbf{0}$ ;*
- (iii)  *$\text{Argmax}(a, x) = [d]_0$  if and only if  $|a_0| = |a_1| + |x_1| = \dots = |a_d| + |x_d| \neq \mathbf{0}$ .*

For this subsection, we fix a TC-hemispace  $G \subseteq \mathbb{T}_{\pm}^d$  with  $G \neq \{\emptyset, \mathbb{T}_{\pm}^d\}$  and let  $(a_0, \dots, a_d) \in \mathbb{T}_{\pm}^{d+1}$ ,  $(a_1, \dots, a_d) \neq \mathbf{0}$ , such that  $\mathcal{H}^+(a) \subseteq G \subseteq \overline{\mathcal{H}^+(a)}$ , which exists by Theorem 5.10. Furthermore, we fix a point  $x \in G$  with  $\text{Argmax}(a, x) \neq$

FIGURE 7.  $\text{Argmax}$  and  $\text{Argmax}^+$  for  $(\ominus 0, \ominus -1, -1)$ 

$\emptyset$ . Observe that we have  $\text{Argmax}^+(a, x) \neq \emptyset$ . Indeed, if  $\text{Argmax}(a, x) \neq \emptyset$  and  $\text{Argmax}^+(a, x) = \emptyset$ , then  $x \in \mathcal{H}^-(a)$ , giving a contradiction with  $x \in G$ .

We start with a statement that be seen orthantwise.

**Lemma 5.13.** *For  $\rho \in \mathbb{R}$  with  $\text{Argmax}(a, x) = \text{Argmax}(a, \rho \odot x)$ , we have  $\rho \odot x \in G$ .*

*Proof.* The claim is trivial for  $\rho = 0$ , so we start with  $\rho < 0$ . If  $0 \in \text{Argmax}(a, x)$ , then the equality  $\text{Argmax}(a, x) = \text{Argmax}(a, \rho \odot x)$  implies that  $\text{Argmax}(a, \rho \odot x) = \{0\} = \text{Argmax}^+(a, \rho \odot x)$  and so  $\rho \odot x \in \mathcal{H}^+(a) \subseteq G$ . If  $0 \notin \text{Argmax}(a, x)$ , then we fix  $k \in \text{Argmax}^+(a, x) \cap [d] = \text{Argmax}^+(a, x)$ . We define a point  $y \in \mathbb{T}_{\pm}^d$  as  $y = \rho \odot x_k \odot e_k$  where  $e_k$  is the  $k$ -th tropical unit vector. We have  $y \in \mathcal{H}^+(a)$  and so  $y \in G$ . Furthermore,  $\rho \odot x = \rho \odot x \triangleleft y$  and therefore  $\rho \odot x \in G$  by the TC-convexity of  $G$ .

If  $\rho > 0$ , then we define  $x' = \rho \odot x$ ,  $G' = \mathbb{T}_{\pm}^d \setminus G$ , and  $a' = \ominus a$ . Suppose that  $x' \in G'$ . Then,  $G'$  is a TC-hemisphere such that  $\mathcal{H}^+(a') \subseteq G' \subseteq \overline{\mathcal{H}^+(a')}$  and we

have  $x = (-\rho) \odot x'$ . Hence, the same reasoning as above shows that  $x \in G'$ , giving a contradiction.  $\square$

**Lemma 5.14.** *For  $y \in \mathbb{T}_{\pm}^d$  with  $\text{Argmax}(a, y) = \text{Argmax}(a, x)$  and  $\text{tsgn}(y_k) = \text{tsgn}(x_k)$  for all  $k \in \text{Argmax}(a, x) \cap [d]$ , we have  $y \in G$ .*

*Proof.* As noted above,  $\text{Argmax}^+(a, x) \neq \emptyset$ , so we fix  $k \in \text{Argmax}^+(a, x)$  and consider the following two cases.

**Case 1** ( $y_j = x_j$  for all  $j \in \text{Argmax}(a, x) \cap [d]$ .)

**Case 1a** ( $|x_j| \leq |y_j|$  for all  $j \notin \text{Argmax}(a, x)$ .)

Define  $M = 0$  if  $k = 0$  and  $M = 1$  otherwise. Let  $z \in \mathbb{T}_{\pm}^d$  be the point defined as

$$z_i = \begin{cases} \mathbf{0} & \text{if } i \in \text{Argmax}(a, y) \setminus \{k\}, \\ M \odot y_i & \text{otherwise.} \end{cases}$$

Then,  $z$  satisfies  $\text{Argmax}(a, z) = \text{Argmax}^+(a, z) = \{k\}$  as one sees by taking scalar product with  $a$ . In particular, we have  $z \in \mathcal{H}^+(a) \subseteq G$ . Moreover,  $|x_j| \leq |y_j|$  for all  $j \notin \text{Argmax}(a, x)$  implies that  $y = (-M) \odot z \triangleleft x$ , and so  $y \in G$ .

**Case 1b** (There exists an  $\ell \notin \text{Argmax}(a, x)$  such that  $|x_\ell| > |y_\ell|$ .)

Let  $u \in \mathbb{T}_{\pm}^d$  be any point that satisfies  $u_j = x_j$  for all  $j \in \text{Argmax}(a, x)$  and  $|u_j| = |x_j|$  otherwise. Then, we have  $\text{Argmax}(a, u) = \text{Argmax}(a, x)$  and  $u \in G$  by the previous case. Since  $u$  was arbitrary, Example 3.19 implies that the point  $v \in \mathbb{T}_{\pm}^d$  defined as

$$v_i = \begin{cases} x_i & \text{if } i \in \text{Argmax}(a, x), \\ \mathbf{0} & \text{otherwise} \end{cases}$$

belongs to  $G$ . Now we can apply the Case 1a with  $v$  instead of  $x$  since  $\mathbf{0} = |v_j| \leq |y_j|$  for all  $j \notin \text{Argmax}(a, x)$ . Hence, we obtain  $y \in G$ .

**Case 2** (There exists an  $\ell \in \text{Argmax}(a, x) \cap [d]$  such that  $y_\ell \neq x_\ell$ .)

Let  $\rho = |y_\ell| \odot |x_\ell|^{\odot -1} \in \mathbb{R}$  and consider the point  $x' = \rho \odot x$ . Then, for each  $j \in \text{Argmax}(a, x) \cap [d]$  we have

$$|a_j| \odot |x'_j| = \rho \odot |a_j| \odot |x_j| = \rho \odot |a_\ell| \odot |x_\ell| = |a_\ell| \odot |y_\ell| = |a_j| \odot |y_j|. \quad (31)$$

This yields  $x'_j = y_j$  for  $j \in \text{Argmax}(a, x) \cap [d]$  because  $\text{tsgn}(x_j) = \text{tsgn}(y_j)$ .

By the case assumption, we have  $\max_{k \in [d]_0} |a_k \odot x_k| \neq \max_{k \in [d]_0} |a_k \odot y_k|$  as  $y_\ell \neq x_\ell$  but  $\text{Argmax}(a, y) = \text{Argmax}(a, x)$ . In particular,  $0 \notin \text{Argmax}(a, y)$ . Furthermore, (31) gives  $|a_0| < |a_\ell| \odot |y_\ell| = |a_\ell| \odot |x'_\ell|$ , so  $0 \notin \text{Argmax}(a, x')$ . This implies that  $0 \notin \text{Argmax}(a, x) \cup \text{Argmax}(a, x')$ . Therefore, we have  $\text{Argmax}(a, x') = \text{Argmax}(a, x)$  by Corollary 5.12 and Lemma 5.13 implies  $x' \in G$ . Since  $x'_j = y_j$  for all  $j \in \text{Argmax}(a, x) \cap [d]$ , Case 1 shows that  $y \in G$ .  $\square$

The structure of the boundary of TC-hemispace relies heavily on the support of the halfspaces sandwiching it. We discuss this in a simple example which is visualized in Fig. 8.

*Example 5.15.* Suppose that  $G \subset \mathbb{T}_{\pm}^2$  is a TC-hemispace such that

$$\{x \in \mathbb{T}_{\pm}^2 \mid x_1 > 0\} \subseteq G \subseteq \{x \in \mathbb{T}_{\pm}^2 \mid x_1 \geq 0\}. \quad (32)$$

Furthermore, suppose that there is a point  $x \in G$  such that  $x_1 = 0$  and let  $y \in \mathbb{T}_{\pm}^2$  be any point such that  $y_1 = 0$ . Then, for  $a = (\ominus 0, 0, \mathbf{0})$  we get  $\text{Argmax}(a, x) = \{0, 1\} = \text{Argmax}(a, y)$  and  $\text{tsgn}(x_1) = \text{tsgn}(y_1)$ , so Lemma 5.14 implies that  $y \in G$ . Hence, there are only two TC-hemispace that satisfy (32), namely  $G =$

$\{x \in \mathbb{T}_{\pm}^2 \mid x_1 > \mathbf{0}\}$  and  $G = \{x \in \mathbb{T}_{\pm}^2 \mid x_1 \geq \mathbf{0}\}$ . By contrast, suppose that  $G' \subset \mathbb{T}_{\pm}^2$  is a TC-hemisphere such that

$$\{x \in \mathbb{T}_{\pm}^2 \mid x_1 > \mathbf{0}\} \subseteq G' \subseteq \{x \in \mathbb{T}_{\pm}^2 \mid x_1 \geq \mathbf{0}\}. \quad (33)$$

Furthermore, suppose that there is a point  $x' \in G'$  such that  $x'_1 = \mathbf{0}$  and let  $y \in \mathbb{T}_{\pm}^2$  be any point such that  $y_1 = \mathbf{0}$ . Then, for  $a = (\mathbf{0}, \mathbf{0}, \mathbf{0})$  we get  $\text{Argmax}(a, x) = \emptyset$  so the assumption of Lemma 5.14 is not satisfied and we cannot deduce anything about  $y$  using this lemma. Indeed, if  $y_2 < x'_2$ , then

$$G' = \{x \in \mathbb{T}_{\pm}^2 \mid x_1 > \mathbf{0}\} \cup \{x \in \mathbb{T}_{\pm}^2 \mid x_1 = \mathbf{0}, x_2 \geq x'_2\}$$

is a TC-hemisphere that does not contain  $y$  and

$$G' = \{x \in \mathbb{T}_{\pm}^2 \mid x_1 > \mathbf{0}\} \cup \{x \in \mathbb{T}_{\pm}^2 \mid x_1 = \mathbf{0}, x_2 \leq x'_2\}$$

is a TC-hemisphere that contains  $y$ . In particular, there are infinitely many TC-hemispheres that satisfy (33).




FIGURE 8. Possibilities of boundary of TC-hemisphere sandwiched between  $x_1 \geq \mathbf{0}$  and  $x_1 > \mathbf{0}$  or  $x_1 \geq \mathbf{0}$  and  $x_1 > \mathbf{0}$ .

**Lemma 5.16.** Define a point  $s \in \mathbb{S}^d$  as

$$\forall \ell \in [d], s_\ell = \begin{cases} x_\ell & \text{if } \ell \in \text{Argmax}^+(a, x), \\ \bullet x_\ell & \text{otherwise,} \end{cases}$$

then  $\mathcal{U}(s) \subseteq G$ .

*Proof.* By Example 3.19, it is enough to show that the vertices of  $\mathcal{U}(s)$  belong to  $G$ . Note that the vertices of  $\mathcal{U}(s)$  are precisely the points  $y \in \mathbb{T}_{\pm}^d$  that satisfy  $|y_j| = |x_j|$  for all  $j \in [d]$  and  $\text{Argmax}^+(a, x) \subseteq \text{Argmax}^+(a, y)$ . To show that every such point belongs to  $G$ , we proceed by induction over  $m = |\text{Argmax}^+(a, y) \setminus \text{Argmax}^+(a, x)|$ .

If  $m = 0$ , then we have  $y_j = x_j$  for all  $j \in \text{Argmax}(a, x)$  and so the claim follows from Lemma 5.14.

If  $m > 0$ , there is a  $k \in \text{Argmax}^+(a, y) \setminus \text{Argmax}^+(a, x)$ . Consider the point  $z \in \mathbb{T}_\pm^d$  defined as

$$z_i = \begin{cases} x_i & \text{if } i = k, \\ y_i & \text{otherwise.} \end{cases}$$

Then, we have  $z \in G$  be the induction hypothesis. Furthermore, since  $|a_0| < 1 + |a_k| + |y_k|$ , the point  $w \in \mathbb{T}_\pm^d$  defined as

$$w_i = \begin{cases} 1 \odot y_k & \text{if } i = k, \\ \mathbf{0} & \text{otherwise} \end{cases}$$

belongs to  $\mathcal{H}^+(a)$ . In particular,  $w \in G$ . Since  $y = (-1) \odot w \triangleleft z$ , we get  $y \in G$ .  $\square$

Combining the previous lemmas leads to the following description, when two points have the same  $\text{Argmax}(a, \cdot)$ . It shows that the structure of hemispaces behaves rather well with respect to fixing  $\text{Argmax}$ . In particular, the following statement extends [30, Lemma 3.1].

**Proposition 5.17.** *Let  $y \in \mathbb{T}_\pm^d$  be such that  $\text{Argmax}(a, x) = \text{Argmax}(a, y)$  and  $\text{Argmax}^+(a, x) \subseteq \text{Argmax}^+(a, y)$ . Then, we have  $y \in G$ .*

*Proof.* Consider the point  $z \in \mathbb{T}_\pm^d$  defined as  $z_j = \text{tsgn}(y_j)|x_j|$  for all  $j \in [d]$ . Since  $\text{Argmax}^+(a, x) \subseteq \text{Argmax}^+(a, y)$ , we have  $z_j = x_j$  for all  $j \in \text{Argmax}^+(a, x)$ . With  $|z_j| \leq |x_j|$  for all  $j \in [d]$ , Lemma 5.16 implies  $z \in G$ . Furthermore,  $\text{Argmax}(a, z) = \text{Argmax}(a, x) = \text{Argmax}(a, y)$  and  $\text{tsgn}(z_j) = \text{tsgn}(y_j)$  for all  $j \in \text{Argmax}(a, z) \cap [d]$ . Hence, we have  $y \in G$  by Lemma 5.14.  $\square$

**5.4. Lifts of TC-hemispace.** As a final ingredient for the proof of Theorem 6.1, we relate TC-hemispace with convex lifts. For this, we construct specific lifts in the sense of Section 4 based on the structural insights for TC-hemispace in Section 5 so far.

**Proposition 5.18.** *If  $G \subseteq \mathbb{T}_\pm^d$  is a TC-hemisphere, then  $G = \text{sval}(\mathbf{G})$  for some convex set  $\mathbf{G} \subseteq \mathbb{K}^d$ .*

*Proof.* First, note that the claim is trivial if  $G \in \{\emptyset, \mathbb{T}_\pm^d\}$ . Otherwise, Theorem 5.10 shows that there exists  $a = (a_0, \dots, a_d) \in \mathbb{T}_\pm^{d+1}$ ,  $(a_1, \dots, a_d) \neq \mathbf{0}$  such that  $\mathcal{H}^+(a) \subseteq G \subseteq \overline{\mathcal{H}^+(a)}$ .

We prove the claim by induction over  $d$ . If  $d = 1$ , then the claim follows from the fact that, for any  $a \in \mathbb{T}_\pm$ , each set  $\{\mathbf{x} \in \mathbb{K} \mid \text{sval}(\mathbf{x}) \sim a\}$  with  $\sim \in \{\geq, >, \leq, <\}$  is convex.

If  $d > 1$ , then we let  $X = \{x \in \mathbb{T}_\pm^d \mid \text{Argmax}(a, x) \neq \emptyset\}$  and its complement  $X' = \mathbb{T}_\pm^d \setminus X$ . Note that  $X'$  is essentially the set of points whose support is disjoint from the support of  $a$ . We will construct a lift for the intersection of  $G$  with  $X$  and  $X'$  and finish by taking their convex hull.

We start by showing that there exists a convex set  $\mathbf{G}^\perp \subseteq \mathbb{K}^d$  with  $\text{sval}(\mathbf{G}^\perp) = G \cap X'$ . If  $X' = \emptyset$ , then we set  $\mathbf{G}^\perp = \emptyset$ . Otherwise, let  $K = \{k \in [d] \mid a_k \neq \mathbf{0}\}$  and note that we have  $a_0 = \mathbf{0}$  and  $X' = \{x \in \mathbb{T}_\pm^d \mid x_k = \mathbf{0} \forall k \in K\}$ .

If  $K = [d]$ , then we take  $\mathbf{G}^\perp = \{0\}$  if  $\mathbf{0} \in G$  and  $\mathbf{G}^\perp = \emptyset$  otherwise. Note that  $G \cap X'$  and  $X' \setminus G$  are TC-convex as intersection of TC-convex sets. If  $K \subsetneq [d]$ , then



let  $\pi: \mathbb{T}_{\pm}^d \rightarrow \mathbb{T}_{\pm}^{d-|K|}$  be the projection that forgets the coordinates from  $K$ . As the components in  $K$  of  $G \cap X'$ ,  $X' \setminus G \subseteq X'$  are all  $\mathbf{O}$ , Proposition 3.17 implies that the set  $\pi(G \cap X')$  is a TC-hemisphere in  $\mathbb{T}_{\pm}^{d-|K|}$ . Hence, by the induction hypothesis, we have  $\pi(G \cap X') = \text{sval}(\mathbf{G}')$  for some convex set  $\mathbf{G}' \subseteq \mathbb{K}^{d-|K|}$ . We embed the set  $\mathbf{G}'$  into  $\mathbb{K}^d$  by adding 0 coordinates to every point and this gives a convex set  $\mathbf{G}^{\perp} \subseteq \mathbb{K}^d$  such that  $\text{sval}(\mathbf{G}^{\perp}) = G \cap X'$ .

Next, we show that there exists a convex set  $\mathbf{G}^{\#} \subseteq \mathbb{K}^d$  with  $\text{sval}(\mathbf{G}^{\#}) = G \cap X$ . To construct an appropriate lift with Definition 4.4, let  $J = \{k \in [d] \mid a_k > \mathbf{O}\}$  be such that  $\overline{\mathcal{H}}^+(a)$  is of type  $J$ . Set  $\mathbf{G}^{\#} = \text{conv}(\{\text{li}_J(x) \in \mathbb{K}^d \mid x \in G \cap X'\})$ . Then,  $\mathbf{G}^{\#}$  is convex and right from the definition we have  $G \cap X \subseteq \text{sval}(\mathbf{G}^{\#})$ .

To prove that  $\text{sval}(\mathbf{G}^{\#}) \subseteq G \cap X$ , let  $\mathbf{y} \in \mathbf{G}^{\#}$  be arbitrary and  $y = \text{sval}(\mathbf{y})$ . Then, the Carathéodory theorem in  $\mathbb{K}^d$  shows that

$$\mathbf{y} = \lambda_1 \text{li}_J(y^{(1)}) + \cdots + \lambda_m \text{li}_J(y^{(m)}) \quad (34)$$

for some  $m \leq d+1$ ,  $y^{(1)}, \dots, y^{(m)} \in G \cap X$ ,  $\lambda_1, \dots, \lambda_m > \mathbf{O}$ , and  $\sum_i \lambda_i = 1$ . As we aim to prove that  $y \in G \cap X$ , we will estimate the leading terms of the components in the sum (34). For every  $i \in [m]$  let  $\lambda_i = \text{sval}(\lambda_i) > \mathbf{O}$  and  $\gamma_i = \text{lc}(\lambda_i) > \mathbf{O}$ . We assume that the points  $y^{(1)}, \dots, y^{(m)}$  are labeled such that  $\gamma_1 \geq \cdots \geq \gamma_m$ . To relate  $y$  with the hemisphere property of  $G$ , we set

$$z = \lambda_1 \odot y^{(1)} \triangleleft \cdots \triangleleft \lambda_m \odot y^{(m)}. \quad (35)$$

With  $\bigoplus_i \lambda_i = \mathbf{O}$ , the TC-convexity of  $G$  implies  $z \in G$ . Moreover, we also have  $z \in X$ . Indeed, this is trivial if  $a_0 \neq \mathbf{O}$ . Otherwise, because of  $y^{(1)} \in X$ , it has a component  $k \in [d]$  such that  $a_k \odot y_k^{(1)} \neq \mathbf{O}$  and so  $a_k \odot z_k \neq \mathbf{O}$ . Thus,  $z \in G \cap X$ .

We will prove that  $y \in G$  using Lemma 5.16. To do so, suppose that there exists  $k \in \text{Argmax}^+(a, z) \cap [d]$ . If  $a_k > \mathbf{O}$  and  $z_k > \mathbf{O}$ , then we define a vector  $u \in \mathbb{R}^m$  by

$$u_i = \begin{cases} \text{lc}\left(\left(\text{li}_J(y^{(i)})\right)_k\right) & \text{if } \left|\lambda_i \odot y_k^{(i)}\right| = |z_k|, \\ 0 & \text{otherwise.} \end{cases}$$

Since  $k \in J$ , we have  $u \in \{-1, 0, d+1\}^m$ . Moreover, since  $z_k > \mathbf{O}$ , the vector  $u$  is nonzero. Let  $p \in [m]$  be the smallest index such that  $u_p \neq 0$ , which is exactly the entry defining the  $k$ th component in the left sum (35). Hence, we have  $z_k = \lambda_p \odot y_k^{(p)}$ . Now,  $z_k > \mathbf{O}$  implies  $y_k^{(p)} > \mathbf{O}$  and therefore  $u_p = d+1$  by the definition of the lift of type  $J$ .

We are ready to estimate the sum of the leading coefficients of the dominating terms in the sum  $\lambda_1 (\text{li}_J(x_1))_k + \cdots + \lambda_m (\text{li}_J(x_m))_k$ . This is just

$$\begin{aligned} \gamma_1 u_1 + \cdots + \gamma_m u_m &\geq (d+1)\gamma_p - \gamma_{p+1} - \cdots - \gamma_m \\ &\geq (d+1)\gamma_p - (m-p)\gamma_p \geq p\gamma_p > \mathbf{O}. \end{aligned} \quad (36)$$

Therefore, there is no cancellation of the leading terms in (34) which implies  $y_k = \text{sval}(\mathbf{y}_k) = z_k$ . Analogously, we obtain  $y_k = z_k$  if  $a_k < \mathbf{O}$  and  $z_k < \mathbf{O}$ . In particular,  $y_k = z_k$  for every  $k \in \text{Argmax}^+(a, z) \cap [d]$ . Furthermore, for every  $\ell \in [d]$  we have  $\ominus|z_\ell| \leq y_\ell \leq |z_\ell|$  by Lemma 4.2. Therefore,  $y \in G$  by Lemma 5.16.

Moreover, we have  $y \in X$ . Indeed, this is trivial if  $a_0 \neq \mathbf{O}$ . Otherwise,  $\text{Argmax}^+(a, z) \cap [d] \neq \emptyset$  and so  $a_k \odot y_k \neq \mathbf{O}$  for at least one  $k \in [d]$ . Thus, we have  $\text{sval}(\mathbf{G}^{\#}) = G \cap X$ .

To finish the proof, let  $\mathbf{G} = \text{conv}(\mathbf{G}^\perp \cup \mathbf{G}^-) \subseteq \mathbb{K}^d$ . Then,  $\mathbf{G}$  is convex and  $G \subseteq \text{sval}(\mathbf{G})$ . To prove that  $\text{sval}(\mathbf{G}) \subseteq G$ , note that this inclusion is trivial if  $a_0 \neq \mathbf{0}$ . Otherwise, let  $\mathbf{z} \in \mathbf{G}$ . Then, there exist  $\mathbf{x} \in \mathbf{G}^\perp, \mathbf{y} \in \mathbf{G}^-, \boldsymbol{\lambda}, \boldsymbol{\mu} \geq \mathbf{0}, \boldsymbol{\lambda} + \boldsymbol{\mu} = \mathbf{1}$  such that  $\mathbf{z} = \boldsymbol{\lambda}\mathbf{x} + \boldsymbol{\mu}\mathbf{y}$ . Denote  $z = \text{sval}(\mathbf{z}), x = \text{sval}(\mathbf{x}), y = \text{sval}(\mathbf{y}), \lambda = \text{sval}(\boldsymbol{\lambda}), \mu = \text{sval}(\boldsymbol{\mu})$ , and  $w = \lambda \odot x \triangleleft \mu \odot y \in G$ . If  $\lambda = \mathbf{0}$  or  $\mu = \mathbf{0}$ , then we trivially have  $z \in G$ . Otherwise,  $y \in X$  and  $\mu \neq \mathbf{0}$  imply that  $w \in X$ . Furthermore,  $x \in X'$  implies that  $z_k = \mu \odot y_k = w_k$  for all  $k \in K$ . Therefore,  $z \in G$  by Lemma 5.14.  $\square$

## 6. TC-HULL AS INTERSECTION OF CLOSED HALFSPPACES

In Section 5, we prepared the crucial tools for proving the representation of the TC-convex hull of finitely many points as the intersection of its containing closed halfspaces. It relies on the understanding of the structure of TC-hemispaces and a representation of those using insights on their lifts from Section 4. Finally, we arrive at analogs of the Minkowski–Weyl theorem for polyhedra.

**Theorem 6.1.** *For every  $x_1, \dots, x_n \in \mathbb{T}_\pm^d$  we have*

$$\text{conv}_{\text{TC}}(x_1, \dots, x_n) = \bigcap_{a \in \mathbb{T}_\pm^{d+1}} \left\{ \overline{\mathcal{H}}^+(a) \mid \{x_1, \dots, x_n\} \subseteq \overline{\mathcal{H}}^+(a) \right\}.$$

*Proof.* The inclusion  $\subseteq$  follows from Corollary 3.8.

For the reverse direction, Lemma 4.13 shows the equality

$$\begin{aligned} & \bigcap_{a \in \mathbb{T}_\pm^{d+1}} \left\{ \overline{\mathcal{H}}^+(a) \mid \{x_1, \dots, x_n\} \subseteq \overline{\mathcal{H}}^+(a) \right\} \\ &= \bigcap \left\{ \text{sval}(\mathbf{X}) \mid \mathbf{X} \subseteq \mathbb{K}^d \text{ is convex and } \{x_1, \dots, x_n\} \subseteq \text{sval}(\mathbf{X}) \right\} \\ &\subseteq \bigcap \left\{ \text{sval}(\mathbf{X}) \mid \mathbf{X} \subseteq \mathbb{K}^d \text{ convex, } \{x_1, \dots, x_n\} \text{ in TC-hemisphere } \text{sval}(\mathbf{X}) \right\}. \end{aligned}$$

By Proposition 5.18, every TC-hemisphere containing  $\{x_1, \dots, x_n\}$  arises in the latter intersection. Combined with Theorem 5.3, this yields

$$\begin{aligned} & \bigcap \left\{ \text{sval}(\mathbf{X}) \mid \mathbf{X} \subseteq \mathbb{K}^d \text{ convex, } \{x_1, \dots, x_n\} \text{ in TC-hemisphere } \text{sval}(\mathbf{X}) \right\} \\ &= \bigcap \left\{ G \mid G \text{ TC-hemisphere with } \{x_1, \dots, x_n\} \subseteq G \right\} \\ &= \text{conv}_{\text{TC}}(x_1, \dots, x_n). \end{aligned} \quad \square$$

Combining Theorem 6.1 and Lemma 4.13 we obtain the following corollary, which is an analog of [42, Theorem 3.14] for TC-convexity.

**Corollary 6.2.** *For every  $x_1, \dots, x_n \in \mathbb{T}_\pm^d$  we have*

$$\begin{aligned} \text{conv}_{\text{TC}}(x_1, \dots, x_n) &= \bigcap \left\{ \text{sval}(\text{conv}(\mathbf{x}_1, \dots, \mathbf{x}_m)) \mid \forall i, \mathbf{x}_i \in \text{sval}^{-1}(x_i) \right\} \\ &= \bigcap_{J \subseteq [d]} \text{sval}(\text{conv}(\text{li}_J(x_1), \dots, \text{li}_J(x_n))). \end{aligned}$$

Based on the description of the TC-convex hull by closed halfspaces, we obtain several further insights on TC-convex hulls. We start with a conic version.

**Theorem 6.3.** *Let  $X \subseteq \mathbb{T}_\pm^d$  be an arbitrary set. Then, the following are equivalent:*

- (i)  $X$  is an intersection of finitely many closed linear tropical halfspaces.
- (ii)  $X = \text{cone}_{\text{TC}}(x_1, \dots, x_m)$  for a finite collection of points  $\{x_1, \dots, x_m\}$ .

(iii) There exists a finite collection of polyhedral cones  $\mathbf{P}_1, \dots, \mathbf{P}_m \subseteq \mathbb{K}^d$  such that

$$\bigcap_i \text{sval}(\mathbf{P}_i) = X.$$

(iv) There exists a finite collection of linear halfspaces  $\mathbf{H}_1, \dots, \mathbf{H}_m \subseteq \mathbb{K}^d$  such that

$$\bigcap_i \text{sval}(\mathbf{H}_i) = X.$$

*Proof.* To prove the implication (i)  $\Rightarrow$  (iv), suppose that  $X = \bigcap_{i=1}^m \overline{\mathcal{H}}^+(a_i)$  for some  $a_1, \dots, a_m \in \mathbb{T}_{\pm}^{d+1}$ . For every  $i \in [m]$ , let  $\mathbf{H}_i = \overline{\mathcal{H}}^+(\text{cli}(a_i))$ . By Lemma 4.6 we have  $X = \bigcap_{i=1}^m \overline{\mathcal{H}}^+(a_i) = \bigcap_{i=1}^m \text{sval}(\mathbf{H}_i)$ .

The implication (iv)  $\Rightarrow$  (iii) is trivial and the implication (iii)  $\Rightarrow$  (i) follows from Lemma 4.14.

To prove the implication (i)  $\Rightarrow$  (ii), we use the tropical Minkowski–Weyl theorem for unsigned tropical convexity [31, Theorem 1]. This theorem implies that for every closed orthant  $O$  of  $\mathbb{T}_{\pm}^d$  there exists a finite set  $U_O \subset O$  such that  $X \cap O = \text{cone}_{\text{TC}}(U_O)$ . Let  $U = \bigcup_O U_O$ . We will show that  $X = \text{cone}_{\text{TC}}(U)$ . Since  $\text{cone}_{\text{TC}}(U_O) \subseteq \text{cone}_{\text{TC}}(U)$  for every  $O$ , we get  $X \subseteq \text{cone}_{\text{TC}}(U)$ . To prove the opposite inclusion, note that  $U \subseteq X$  implies  $\text{conv}_{\text{TC}}(U) \subseteq X$  by Corollary 3.8. Since  $X$  is an intersection of linear tropical halfspaces, we get  $\text{cone}_{\text{TC}}(U) \subseteq X$ .

To finish the proof, it is enough to show the implication (ii)  $\Rightarrow$  (iii). To do so, let

$$\mathbf{P}_J = \text{cone}(\text{li}_J(x_1), \dots, \text{li}_J(x_m))$$

for all  $J \subseteq [d]$ . We will show that  $X = \bigcap_{J \subseteq [d]} \text{sval}(\mathbf{P}_J)$ . If  $x \in X$ , there exists  $\lambda \in \mathbb{T}_{\geq 0}$  such that  $x \in \lambda \odot \text{conv}_{\text{TC}}(x_1, \dots, x_m)$ . By combining Theorem 6.1 with Lemma 4.13, for all  $J \subseteq [d]$ , we get

$$\text{conv}_{\text{TC}}(x_1, \dots, x_m) \subseteq \text{sval}\left(\text{conv}(\text{li}_J(x_1), \dots, \text{li}_J(x_m))\right).$$

Hence, by Lemma 4.15 we have

$$\begin{aligned} \lambda \odot \text{conv}_{\text{TC}}(x_1, \dots, x_m) &\subseteq \lambda \odot \text{sval}\left(\text{conv}(\text{li}_J(x_1), \dots, \text{li}_J(x_m))\right) \\ &= \text{sval}\left(\text{conv}(t^\lambda \text{li}_J(x_1), \dots, t^\lambda \text{li}_J(x_m))\right) \subseteq \text{sval}(\mathbf{P}_J) \end{aligned}$$

and  $x \in \bigcap_{J \subseteq [d]} \text{sval}(\mathbf{P}_J)$ . Conversely, if  $x \in \bigcap_{J \subseteq [d]} \text{sval}(\mathbf{P}_J)$  then, by Lemma 4.15, for every  $J \subseteq [d]$  there exists  $\lambda_J \in \mathbb{T}_{\geq 0}$  such that

$$\begin{aligned} x &\in \text{sval}\left(\text{conv}(t^{\lambda_J} \text{li}_J(x_1), \dots, t^{\lambda_J} \text{li}_J(x_m))\right) \\ &= \text{sval}\left(\text{conv}(\text{li}_J(\lambda_J \odot x_1), \dots, \text{li}_J(\lambda_J \odot x_m))\right). \end{aligned}$$

Hence, by combining Theorem 6.1 with Lemma 4.13 and Lemma 3.30 we get

$$\begin{aligned} x &\in \bigcap_J \text{sval}\left(\text{conv}\left\{\text{li}_J(\lambda_J \odot x_i) \mid \tilde{J} \subseteq [d], i \in [m]\right\}\right) \\ &= \text{conv}_{\text{TC}}\left(\left\{\lambda_J \odot x_i \mid \tilde{J} \subseteq [d], i \in [m]\right\}\right) \\ &\subseteq \text{cone}_{\text{TC}}(x_1, \dots, x_m) = X. \end{aligned} \quad \square$$

Establishing an affine version of the former theorem requires us to come up with an appropriate concept of dehomogenization.

**Theorem 6.4.** *Let  $X \subseteq \mathbb{T}_{\pm}^d$  be an arbitrary set. Then, the following are equivalent:*

- (i)  $X$  is an intersection of finitely many closed tropical halfspaces.
- (ii) There exist two finite sets  $V, W \subset \mathbb{T}_{\pm}^d$  such that

$$\text{conv}_{\text{TC}}(\{v \triangleleft \lambda \odot w \mid v \in V, w \in W, \lambda \in \mathbb{T}_{\geq \mathbf{0}}\}) = X.$$

- (iii) There exists a finite collection of polyhedra  $\mathbf{P}_1, \dots, \mathbf{P}_m \subset \mathbb{K}^d$  such that

$$\bigcap_i \text{sval}(\mathbf{P}_i) = X.$$

- (iv) There exists a finite collection of affine halfspaces  $\mathbf{H}_1, \dots, \mathbf{H}_m \subseteq \mathbb{K}^d$  such that

$$\bigcap_i \text{sval}(\mathbf{H}_i) = X.$$

*Proof.* The theorem is trivial if  $X$  is empty. From now on we suppose that  $X$  is nonempty.

The implication (i)  $\Rightarrow$  (iv) follows by the same argument as in the proof of Theorem 6.3.

The implication (iv)  $\Rightarrow$  (iii) is trivial.

To prove that (iii)  $\Rightarrow$  (ii), for every  $i \in [m]$ , let  $\mathbf{Q}_i = \{(\lambda \mathbf{x}, \lambda) : \lambda \geq 0, \mathbf{x} \in \mathbf{P}_i\} \subseteq \mathbb{K}^{d+1}$ . The set  $\mathbf{Q}_i$  is a polyhedral cone. Furthermore, we have the equality

$$\text{sval}(\mathbf{Q}_i) \cap \{x_{d+1} = 0\} = \{(x, 0) \mid x \in \text{sval}(\mathbf{P}_i)\}. \quad (37)$$

Indeed, if  $x \in \text{sval}(\mathbf{P}_i)$ , then there exists  $\mathbf{x} \in \mathbf{P}_i$  such that  $\text{sval}(\mathbf{x}) = x$ . Therefore,  $(\mathbf{x}, 1) \in \mathbf{Q}_i$  and  $(x, 0) \in \text{sval}(\mathbf{Q}_i)$ . Conversely, if  $(x, 0) \in \text{sval}(\mathbf{Q}_i)$ , then there exist  $\lambda \geq 0$  and  $\mathbf{x} \in \mathbf{P}_i$  such that  $\text{sval}(\lambda) = 0$  and  $x = \text{sval}(\lambda \mathbf{x}) = \text{sval}(\lambda) + \text{sval}(\mathbf{x}) = \text{sval}(\mathbf{x})$ . Therefore  $x \in \text{sval}(\mathbf{P}_i)$  and (37) is satisfied. Let  $Y = \bigcap_{i=1}^m \text{sval}(\mathbf{Q}_i) \subseteq \mathbb{T}_{\pm}^{d+1}$ . By (37), we get

$$Y \cap \{x_{d+1} = 0\} = \{(x, 0) \mid x \in X\}. \quad (38)$$

Applying Theorem 6.3 to the set  $Y$ , there exists a finite set  $U = \{u_1, \dots, u_n\} \subset \mathbb{T}_{\pm}^{d+1}$  such that  $Y = \text{cone}_{\text{TC}}(U)$ . Since  $Y$  is a cone, we can suppose that  $U$  contains  $\mathbf{0}$ . Moreover, we have  $y_{d+1} \geq \mathbf{0}$  for every  $y \in Y$ . Hence, we can scale every point  $u^{(i)} \in U$  in such a way that  $u_{d+1}^{(i)} \in \{\mathbf{0}, 0\}$ . Thus, we can write  $U = \hat{V} \cup \hat{W}$  where  $\hat{V}$  contains the elements of  $U$  whose last coordinate is 0 and  $\hat{W}$  contains the elements whose last coordinate is  $\mathbf{0}$ . Since  $U$  contains  $\mathbf{0}$ , the set  $\hat{W}$  is nonempty. Since  $X$  is nonempty, (38) implies that  $\hat{V}$  is also nonempty. Therefore, by combining (38) with Lemma 3.32 we get

$$X = \text{conv}_{\text{TC}}(\{v \triangleleft \lambda \odot w \mid v \in V, w \in W, \lambda \in \mathbb{T}_{\geq \mathbf{0}}\}),$$

where  $V, W$  are the projections of  $\hat{V}, \hat{W}$  obtained by deleting the last coordinate.

To prove the implication (ii)  $\Rightarrow$  (i), we define  $\hat{V} = \{(v, 0) : v \in V\}$  and  $\hat{W} = \{(w, \mathbf{0}) : w \in \hat{W}\}$ . By Theorem 6.3, we have

$$\text{cone}_{\text{TC}}(\hat{V} \cup \hat{W}) = \bigcap_{i=1}^n \{x \in \mathbb{T}_{\pm}^{d+1} \mid a_{i,1} \odot x_1 \oplus \dots \oplus a_{i,d+1} \odot x_{d+1} \neq \mathbf{0}\}$$

for some finite set  $\{a_1, \dots, a_n\} \in \mathbb{T}_{\pm}^{d+1} \setminus \{\mathbf{0}\}$ . By Lemma 3.32 we have the equality  $\{(x, 0) : x \in X\} = \text{cone}_{\text{TC}}(\hat{V} \cup \hat{W}) \cap \{x_{d+1} = 0\}$  and therefore

$$X = \bigcap_{i=1}^n \left\{ x \in \mathbb{T}_{\pm}^{d+1} \mid a_{i,d+1} \oplus a_{i,1} \odot x_1 \oplus \dots \oplus a_{i,d} \odot x_d \neq \mathbf{0} \right\}. \quad \square$$

We conclude with a strengthening of the separation theorems in [42, Section 5] using the Pash property of TO-convexity shown in Theorem 3.3.

**Theorem 6.5.** *Suppose that a set  $X \subseteq \mathbb{T}_{\pm}^d$  is TO-convex. Then, the set  $\text{cl}(X)$  is equal to the intersection of closed tropical halfspaces that contain it.*

*Furthermore, if  $X$  is a nonempty TO-convex cone, then  $\text{cl}(X)$  is equal to the intersection of linear closed tropical halfspaces that contain it.*

*Proof.* Let  $X$  be a TO-convex set. It is obvious that  $\text{cl}(X)$  is included in the intersection of closed tropical halfspaces that contain it. To prove the opposite inclusion, let  $y \notin X$ . We can find  $\ell_1, \dots, \ell_d, r_1, \dots, r_d \in \mathbb{T}_{\pm}$  such that

$$\ell_1 < y_1 < r_1, \ell_2 < y_2 < r_2, \dots, \ell_d < y_d < r_d$$

and such that the box  $B = [\ell_1, r_1] \times [\ell_2, r_2] \times \dots \times [\ell_d, r_d]$  does not intersect  $\text{cl}(X)$ . Since  $B$  and  $X$  are TO-convex and disjoint, Theorem 3.3 implies that there exists a TO-hemisphere  $G$  such that  $X \subseteq G$  and  $B \subseteq \mathbb{T}_{\pm}^d \setminus G$ . Therefore, by Theorem 5.10, there exists a closed tropical halfspace  $\overline{\mathcal{H}}^+(a)$  such that  $\text{cl}(X) \subseteq \overline{\mathcal{H}}^+(a)$  and  $B \subseteq \overline{\mathcal{H}}^-(a)$ . Furthermore,  $y$  belongs to the interior of  $B$  and so we have  $y \in \mathcal{H}^-(a)$  by Lemma 2.4. Since  $y$  was arbitrary, we get the first claim.

To prove the second claim, suppose that  $X$  is a nonempty TO-convex cone and let  $y \notin \text{cl}(X)$ . By the first part of the theorem, there exists  $a \in \mathbb{T}_{\pm}^{d+1}$  such that  $\text{cl}(X) \subseteq \overline{\mathcal{H}}^+(a)$  and  $y \in \mathcal{H}^-(a)$ . Since  $X$  is a nonempty cone, we have  $\mathbf{0} \in \text{cl}(X)$ . In particular,  $a_0 \geq \mathbf{0}$ . Let  $\tilde{a} = (\mathbf{0}, a_1, \dots, a_d)$ . Then,  $y \in \mathcal{H}^-(\tilde{a})$ . Suppose that  $x \in \text{cl}(X)$  is such that  $x \notin \overline{\mathcal{H}}^+(\tilde{a})$ . Then,  $a_1 \odot x_1 \oplus \dots \oplus a_d \odot x_d < \mathbf{0}$  and so  $\lambda \odot x \in \mathcal{H}^-(a)$  for a sufficiently large  $\lambda > \mathbf{0}$ . This gives a contradiction with  $\text{cl}(X) \subseteq \overline{\mathcal{H}}^+(a)$ . Hence, we have  $\text{cl}(X) \subseteq \overline{\mathcal{H}}^+(\tilde{a})$ , which finishes the proof.  $\square$

## 7. CONCLUSION

One of our main results is the representation of a finitely generated TC-convex set as an intersection of closed tropical halfspaces in Theorem 6.1. Furthermore, we show in Theorem 4.10 that also the valuation of a closed convex semialgebraic set has such a representation. This motivates the following.

*Conjecture 7.1.* For all closed TC-convex sets, we have

$$X = \bigcap_{a \in \mathbb{T}_{\pm}^{d+1}} \left\{ \overline{\mathcal{H}}^+(a) \mid X \subseteq \overline{\mathcal{H}}^+(a) \right\}. \quad (39)$$

Such a more general statement could be deduced from a more direct proof of the representation by halfspaces. On one hand, it would be interesting to get stronger separation without relying on the separation results over Puiseux series by using Lemma 4.13 in the proof of Theorem 6.1. On the other hand, the proof of Theorem 5.3 uses the Kuratowski–Zorn lemma which is highly non-constructive.

*Question 7.2.* How can one deduce Theorem 6.1 in a more constructive way?

Example 3.28 discusses the Carathéodory number  $c_d$  of TC-convexity. The examples give a lower bound  $c_d \geq 2^d$  complementing the upper bound  $c_d \leq d2^d + 1$  given in Proposition 3.27. Recall that TC-convexity extends the ‘usual’ tropical convexity which has Carathéodory number  $d + 1$  as already shown in [34, 28, 21].

*Question 7.3.* What is the Carathéodory number of TC-convexity?

An approach for the former question would be via a better understanding of the operator  $\text{Faces}(\cdot)$ . In Definition 3.11, we introduced it as a crucial building block for the structure of TC-convex sets. On a more abstract level, the operator produces certain cubical subcomplexes of the cubical complex formed by the faces of a hypercube. A better understanding of these complexes might lead to a better bound.

#### ACKNOWLEDGEMENTS

We thank the 2021 HIM program Discrete Optimization during which part of this work was developed.

#### REFERENCES

- [1] M. Akian, G. Cohen, S. Gaubert, R. Nikoukhah, and J. P. Quadrat. Linear systems in  $(\max, +)$  algebra. In *29th IEEE Conference on Decision and Control*, pages 151–156 vol.1, Dec 1990.
- [2] Marianne Akian, Stéphane Gaubert, and Alexander Guterman. Tropical polyhedra are equivalent to mean payoff games. *Internat. J. Algebra Comput.*, 22(1):1250001, 43, 2012.
- [3] Marianne Akian, Stéphane Gaubert, and Alexander Guterman. Tropical Cramer determinants revisited. In *Tropical and idempotent mathematics and applications*, volume 616 of *Contemp. Math.*, pages 1–45. Amer. Math. Soc., Providence, RI, 2014.
- [4] Daniele Alessandrini. Logarithmic limit sets of real semi-algebraic sets. *Adv. Geom.*, 13(1):155–190, 2013.
- [5] X. Allamigeon, P. Benchimol, S. Gaubert, and M. Joswig. Combinatorial simplex algorithms can solve mean payoff games. *SIAM J. Optim.*, 24(4):2096–2117, 2014.
- [6] X. Allamigeon, S. Gaubert, R. D. Katz, and M. Skomra. Universal complexity bounds based on value iteration and application to entropy games. In *49th International Colloquium on Automata, Languages, and Programming (ICALP 2022)*, volume 229 of *Leibniz International Proceedings in Informatics (LIPIcs)*, pages 126:1–126:20, Dagstuhl, Germany, 2022. Schloss Dagstuhl – Leibniz-Zentrum für Informatik. To appear.
- [7] X. Allamigeon, S. Gaubert, and M. Skomra. Solving generic nonarchimedean semidefinite programs using stochastic game algorithms. *J. Symbolic Comput.*, 85:25–54, 2018.
- [8] X. Allamigeon, S. Gaubert, and M. Skomra. The tropical analogue of the Helton–Nie conjecture is true. *J. Symbolic Comput.*, 91:129–148, 2019.
- [9] Xavier Allamigeon, Pascal Benchimol, Stéphane Gaubert, and Michael Joswig. Tropicalizing the simplex algorithm. *SIAM J. Discrete Math.*, 29(2):751–795, 2015.
- [10] Xavier Allamigeon, Pascal Benchimol, Stéphane Gaubert, and Michael Joswig. Log-barrier interior point methods are not strongly polynomial. *SIAM J. Appl. Algebra Geom.*, 2(1):140–178, 2018.
- [11] Xavier Allamigeon, Pascal Benchimol, Stéphane Gaubert, and Michael Joswig. What tropical geometry tells us about the complexity of linear programming. *SIAM Review*, 63(1):123–164, January 2021.
- [12] Xavier Allamigeon, Stéphane Gaubert, and Ricardo D. Katz. Tropical polar cones, hypergraph transversals, and mean payoff games. *Linear Algebra Appl.*, 435(7):1549–1574, 2011.
- [13] Xavier Allamigeon, Stéphane Gaubert, and Mateusz Skomra. Tropical spectrahedra. *Discrete Comput. Geom.*, 63:507–548, 2020.
- [14] Xavier Allamigeon, Stéphane Gaubert, and Nicolas Vandame. No self-concordant barrier interior point method is strongly polynomial. In *Proceedings of the 54th Annual ACM SIGACT*

- Symposium on Theory of Computing*, STOC 2022, page 515–528, New York, NY, USA, 2022. Association for Computing Machinery.
- [15] Laura Anderson. Vectors of matroids over tracts. *J. Comb. Theory, Ser. A*, 161:236–270, 2019.
- [16] Matthias Aschenbrenner and Andreas Fischer. Definable versions of theorems by Kirszbraun and Helly. *Proceedings of the London Mathematical Society*, 102(3):468–502, 2011.
- [17] M. Baker and N. Bowler. Matroids over partial hyperstructures. *Adv. Math.*, 343:821–863, 2019.
- [18] S. Basu, R. Pollack, and M.-F. Roy. *Algorithms in Real Algebraic Geometry*, volume 10 of *Algorithms Comput. Math.* Springer, Berlin, 2006.
- [19] Anders Björner, Michel Las Vergnas, Bernd Sturmfels, Neil White, and Günter M. Ziegler. *Oriented matroids*, volume 46 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, 1993.
- [20] Walter Briec. Some remarks on an idempotent and non-associative convex structure. *J. Convex Anal.*, 22(1):259–289, 2015.
- [21] Walter Briec and Charles Horvath.  $\mathbb{B}$ -convexity. *Optimization*, 53(2):103–127, 2004.
- [22] Walter Briec and Charles Horvath. Halfspaces and Hahn–Banach like properties in  $\mathbb{B}$ -convexity and max-plus convexity. *Pac. J. Optim.*, 4(2):293–317, 2008.
- [23] Marcel Celaya. *Lattice points, oriented matroids, and zonotopes*. PhD thesis, Georgia Institute of Technology, 2019.
- [24] Victor Chepoi. Separation of two convex sets in convexity structures. *Journal of Geometry*, 50:30–51, 1994.
- [25] Guy Cohen, Stéphane Gaubert, Jean-Pierre Quadrat, and Ivan Singer. Max-plus convex sets and functions. In *Idempotent mathematics and mathematical physics*, volume 377 of *Contemp. Math.*, pages 105–129. Amer. Math. Soc., Providence, RI, 2005.
- [26] Thomas Colcombet, Nathanaël Fijalkow, Paweł Gawrychowski, and Pierre Ohlmann. The theory of universal graphs for infinite duration games, 2021.
- [27] A. Connes and C. Consani. The hyperring of adèle classes. *J. Number Theory*, 131(2):159–194, 2011.
- [28] Mike Develin and Bernd Sturmfels. Tropical convexity. *Doc. Math.*, 9:1–27, 2004.
- [29] Mike Develin and Josephine Yu. Tropical polytopes and cellular resolutions. *Experiment. Math.*, 16(3):277–291, 2007.
- [30] Daniel Ehrmann, Zach Higgins, and Viorel Nitica. The geometric structure of max-plus hemispaces. *Linear Algebra and its Applications*, 489:144 – 166, 2016.
- [31] Stéphane Gaubert and Ricardo D. Katz. Minimal half-spaces and external representation of tropical polyhedra. *J. Algebraic Combin.*, 33(3):325–348, 2011.
- [32] Stéphane Gaubert and Frédéric Meunier. Carathéodory, Helly and the others in the max-plus world. *Discrete Comput. Geom.*, 43(3):648–662, 2010.
- [33] Stéphane Gaubert and Sergeï Sergeev. Cyclic projectors and separation theorems in idempotent convex geometry. *J. Math. Sci.*, 155:815–829, 2008.
- [34] Siegfried Helbig. On Carathéodory’s and Kreĭn–Milman’s theorems in fully ordered groups. *Comment. Math. Univ. Carolin.*, 29(1):157–167, 1988.
- [35] Horvath, Charles. Some general principles in tropical convexities. *ESAIM: ProcS*, 57:48–63, 2017.
- [36] Philipp Jell, Claus Scheiderer, and Josephine Yu. Real Tropicalization and Analytification of Semialgebraic Sets. *International Mathematics Research Notices*, 2022(2):928–958, 05 2020.
- [37] M. Joswig. *Essentials of tropical combinatorics*, volume 219 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2022.
- [38] Michael Joswig and Georg Loho. Weighted digraphs and tropical cones. *Linear Algebra Appl.*, 501:304–343, 2016.
- [39] Ricardo D. Katz, Viorel Nitica, and Sergeï Sergeev. Characterization of tropical hemispaces by  $(p,r)$ -decompositions. *Linear Algebra and its Applications*, 440:131 – 163, 2014.
- [40] Wiesław Kubiś. *Abstract convex structures in topology and set theory*. PhD thesis, University of Silesia, 1999.
- [41] Wiesław Kubiś. Separation properties of convexity spaces. *Journal of Geometry*, 74:110–119, 2002.
- [42] Georg Loho and László A. Végh. Signed Tropical Convexity. In Thomas Vidick, editor, *11th Innovations in Theoretical Computer Science Conference (ITCS 2020)*, volume 151 of *Leibniz*

- International Proceedings in Informatics (LIPIcs)*, pages 24:1–24:35, Dagstuhl, Germany, 2020. Schloss Dagstuhl–Leibniz-Zentrum fuer Informatik.
- [43] Petros Maragos, Vasileios Charisopoulos, and Emmanouil Theodosis. Tropical geometry and machine learning. *Proceedings of the IEEE*, 109:728–755, 2021.
- [44] D. Marker. *Model Theory: An Introduction*, volume 217 of *Grad. Texts in Math.* Springer, New York, 2002.
- [45] T. Markwig. A field of generalised Puiseux series for tropical geometry. *Rend. Semin. Mat., Univ. Politec. Torino*, 68(1):79–92, 2010.
- [46] R. O. Robson. Separating points from closed convex sets over ordered fields and a metric for  $\tilde{\mathbb{R}}^n$ . *Trans. Amer. Math. Soc.*, 326(1):89–99, 1991.
- [47] R. T. Rockafellar. *Convex Analysis*. Princeton University Press, Princeton, NJ, 1972.
- [48] S. N. Samborskii and G. B. Shpiz. Convex sets in the semimodule of bounded functions. In V. P. Maslov and S. N. Samborskii, editors, *Idempotent analysis*, volume 13 of *Adv. Soviet Math.*, pages 135–137. AMS, Providence, RI, 1992.
- [49] A. Schrijver. *Theory of linear and integer programming*. Wiley-Intersci. Ser. Discrete Math. Optim. Wiley, New York, 1987.
- [50] Alfred Tarski. *A Decision Method for Elementary Algebra and Geometry*. RAND Corporation, Santa Monica, Calif., 1948.
- [51] M. L. J. van de Vel. *Theory of Convex Structures*. North-Holland, Amsterdam, 1993.
- [52] L. van den Dries and P. Speissegger. The real field with convergent generalized power series. *Trans. Amer. Math. Soc.*, 350(11):4377–4421, 1998.
- [53] Oleg Viro. Hyperfields for tropical geometry i. hyperfields and dequantization, 2010.
- [54] Ruriko Yoshida, Leon Zhang, and Xu Zhang. Tropical principal component analysis and its application to phylogenetics. *Bulletin of Mathematical Biology*, 81(2):568–597, Feb 2019.
- [55] Josephine Yu. Tropicalizing the positive semidefinite cone. *Proc. Amer. Math. Soc.*, 143(5):1891–1895, 2015.
- [56] Karel Zimmermann. A general separation theorem in extremal algebras. *Ekonom.-Mat. Obzor*, 13(2):179–201, 1977.
- [57] Uri Zwick and Mike Paterson. The complexity of mean payoff games on graphs. *Theoret. Comput. Sci.*, 158(1–2):343–359, 1996.

#### APPENDIX A. DETAILS ON SEMIALGEBRAIC SETS

We give more explanations for the statement and proof of Proposition 4.8. We refer to [44, Corollary 3.3.20] for another example of a proof that works in the same way.

*Extended proof of Proposition 4.8.* Recall that  $\mathbf{X}, \mathbf{Y} \subseteq \mathbb{K}^d$  are two nonempty convex semialgebraic sets such that  $\mathbf{X} \cap \mathbf{Y} = \emptyset$ . Since  $\mathbf{X}, \mathbf{Y}$  are semialgebraic, there exist two first-order formulas  $\phi(x_1, \dots, x_d, y_1, \dots, y_{n_1}), \psi(x_1, \dots, x_d, y_1, \dots, y_{n_2})$  in the language of ordered fields and two vectors  $\mathbf{r} \in \mathbb{K}^{n_1}, \mathbf{s} \in \mathbb{K}^{n_2}$  such that

$$\begin{aligned}\mathbf{X} &= \{\mathbf{x} \in \mathbb{K}^d : \phi(\mathbf{x}, \mathbf{r}) \text{ is true in } \mathbb{K}\}, \\ \mathbf{Y} &= \{\mathbf{x} \in \mathbb{K}^d : \psi(\mathbf{x}, \mathbf{s}) \text{ is true in } \mathbb{K}\}.\end{aligned}$$

For every  $\mathbf{w} \in \mathbb{K}^{n_1}, \mathbf{z} \in \mathbb{K}^{n_2}$ , let  $\mathbf{X}_{\mathbf{w}}, \mathbf{Y}_{\mathbf{z}}$  be the semialgebraic sets defined as

$$\begin{aligned}\mathbf{X}_{\mathbf{w}} &= \{\mathbf{x} \in \mathbb{K}^d : \phi(\mathbf{x}, \mathbf{w}) \text{ is true in } \mathbb{K}\}, \\ \mathbf{Y}_{\mathbf{z}} &= \{\mathbf{x} \in \mathbb{K}^d : \psi(\mathbf{x}, \mathbf{z}) \text{ is true in } \mathbb{K}\}.\end{aligned}$$

Now, the statement

“For every  $\mathbf{w}, \mathbf{z}$  such that  $\mathbf{X}_{\mathbf{w}}, \mathbf{Y}_{\mathbf{z}}$  are nonempty, convex, and disjoint, there exists a vector  $\mathbf{a}$  such that  $\mathbf{X}_{\mathbf{w}} \subseteq \overline{\mathcal{H}}^+(\mathbf{a})$  and  $\mathbf{Y}_{\mathbf{z}} \subseteq \overline{\mathcal{H}}^-(\mathbf{a})$ ”



can be written as sentence in the language of ordered fields. This sentence is true over  $\mathbb{R}$  by the hyperplane separation theorem, and so it is true over  $\mathbb{K}$  by the completeness of the theory of real closed fields. By taking  $(\mathbf{w}, \mathbf{z}) = (\mathbf{r}, \mathbf{s})$  we obtain the claim.  $\square$

*Example A.1.* The assumption that both sets  $\mathbf{X}, \mathbf{Y}$  are semialgebraic cannot be dropped from Proposition 4.8 as the following example, taken from [46], shows. Let  $\mathbf{A} = \{z \in \mathbb{K} \mid z \geq 0 \wedge \text{val}(z) < 0\}$  and  $\mathbf{B} = \{z \in \mathbb{K} \mid z \geq 0 \wedge \text{val}(z) \geq 0\}$ . In this way,  $\mathbf{A} \cup \mathbf{B}$  is the set of nonnegative Puiseux series,  $\mathbf{A}$  does not have a least upper bound in  $\mathbb{K}$ ,  $\mathbf{B}$  does not have a greatest lower bound in  $\mathbb{K}$ , and  $\mathbf{a} < \mathbf{b}$  for all  $\mathbf{a} \in \mathbf{A}, \mathbf{b} \in \mathbf{B}$ . Let  $\mathbf{X} = \mathbf{X}_1 \cup \mathbf{X}_2$ , where

$$\begin{aligned} \mathbf{X}_1 &= \{(\mathbf{x}, \mathbf{y}) \in \mathbb{K}^2 \mid \mathbf{x} \geq 0 \wedge (\forall z \in \mathbf{B}, \mathbf{y} \leq z\mathbf{x})\} . \\ \mathbf{X}_2 &= \{(\mathbf{x}, \mathbf{y}) \in \mathbb{K}^2 \mid \mathbf{x} \leq 0 \wedge (\forall z \in \mathbf{A}, \mathbf{y} \leq z\mathbf{x})\} . \end{aligned}$$

We note that  $\mathbf{X}$  is nonempty because it contains  $(0, 0)$ . Also, both  $\mathbf{X}_1$  and  $\mathbf{X}_2$  arise as intersections of closed convex sets, so  $\mathbf{X}_1$  and  $\mathbf{X}_2$  are both convex and closed. Hence  $\mathbf{X}$  is closed. To see that  $\mathbf{X}$  is also convex, let  $(\mathbf{x}_1, \mathbf{y}_1) \in \mathbf{X}_1$  and  $(\mathbf{x}_2, \mathbf{y}_2) \in \mathbf{X}_2$  be such that  $\mathbf{x}_1 > 0$  and  $\mathbf{x}_2 < 0$ . Pick  $\lambda \in [0, 1]$  such that  $\lambda\mathbf{x}_1 + (1 - \lambda)\mathbf{x}_2 = 0$  and let  $\mathbf{z} = \max(0, \mathbf{y}_1/\mathbf{x}_1)$ . Since  $\mathbf{z}$  is a lower bound for  $\mathbf{B}$ , we have  $\mathbf{z} \in \mathbf{A}$ . Hence  $\lambda\mathbf{y}_1 + (1 - \lambda)\mathbf{y}_2 \leq \mathbf{z}(\lambda\mathbf{x}_1 + (1 - \lambda)\mathbf{x}_2) = 0$ . Therefore, the point  $\lambda(\mathbf{x}_1, \mathbf{y}_1) + (1 - \lambda)(\mathbf{x}_2, \mathbf{y}_2)$  belongs to both  $\mathbf{X}_1$  and  $\mathbf{X}_2$ . Hence, the whole segment between these two points belongs to  $\mathbf{X}$ .

Despite the fact that  $\mathbf{X}$  is closed and convex, it cannot be separated by a hyperplane from *any* point  $\mathbf{w} \notin \mathbf{X}$ . To see that, suppose that  $(\mathbf{c}_0, \mathbf{c}_1, \mathbf{c}_2)$  is such that  $(\mathbf{c}_1, \mathbf{c}_2) \neq (0, 0)$  and  $\mathbf{c}_0 + \mathbf{c}_1\mathbf{x} + \mathbf{c}_2\mathbf{y} \geq 0$  for all  $(\mathbf{x}, \mathbf{y}) \in \mathbf{X}$ . Since  $(t^x, 0), (0, -t^x), (-t^x, -t^x), (t^x, t^{x-0.5}) \in \mathbf{X}$  for arbitrarily large values of  $x > 0$ , we have  $\mathbf{c}_1 > 0$  and  $\mathbf{c}_2 < 0$ . Let  $\mathbf{z} = \mathbf{c}_1/|\mathbf{c}_2|$ . If  $\mathbf{z} \in \mathbf{A}$ , then by taking any  $\mathbf{z}' \in \mathbf{A}$  that is greater than  $\mathbf{z}$  and considering the points  $(t^x, \mathbf{z}'t^x) \in \mathbf{X}$  we get  $\mathbf{c}_0 + \mathbf{c}_1t^x + \mathbf{c}_2\mathbf{z}'t^x \geq 0 \iff \mathbf{z}' \leq \mathbf{z} + (\mathbf{c}_0/|\mathbf{c}_2|)t^{-x}$ , which is a contradiction for  $x$  large enough. Likewise, if  $\mathbf{z} \in \mathbf{B}$ , then by taking any  $\mathbf{z}' \in \mathbf{B}$  that is smaller than  $\mathbf{z}$  and considering the points  $(-t^x, -\mathbf{z}'t^x) \in \mathbf{X}$  we get  $\mathbf{c}_0 - \mathbf{c}_1t^x - \mathbf{c}_2\mathbf{z}'t^x \geq 0 \iff \mathbf{z}' \geq \mathbf{z} - (\mathbf{c}_0/|\mathbf{c}_2|)t^{-x}$ , which is a contradiction for  $x$  large enough.

(Georg Loho) UNIVERSITY OF TWENTE, DEPARTMENT OF APPLIED MATHEMATICS, THE NETHERLANDS,

*Email address:* g.loho@utwente.nl

(Mateusz Skomra) LAAS-CNRS, UNIVERSITÉ DE TOULOUSE, CNRS, TOULOUSE, FRANCE,

*Email address:* mateusz.skomra@laas.fr