



HAL
open science

A modified Christoffel function and its asymptotic properties

Jean-Bernard Lasserre

► **To cite this version:**

Jean-Bernard Lasserre. A modified Christoffel function and its asymptotic properties. 2023. hal-03949201v1

HAL Id: hal-03949201

<https://laas.hal.science/hal-03949201v1>

Preprint submitted on 23 Jan 2023 (v1), last revised 26 Jan 2023 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

ON A PARTICULAR EXTENSION OF THE CHRISTOFFEL FUNCTION

JBL

1. INTRODUCTION

A certain variant of the Christoffel function (see below) allows to relate to the underlying density without scaling factor. This variant considers balls $\mathbf{B}(\boldsymbol{\xi}, \delta)$ around each point $\boldsymbol{\xi} \in \text{supp}(\mu)$, as also do people in topological data analysis (see the interesting paper [1]).

A preliminary result. Let $\Omega \subset \mathbb{R}^d$ be compact with nonempty interior and such that $\Omega = \overline{\text{int}(\Omega)}$.

Lemma 1.1. *Let $\phi, \psi \in \mathcal{M}(\Omega)_+$ be two finite Borel measures on Ω with $\mathbf{M}_n(\phi), \mathbf{M}_n(\psi) > 0$ for all $n \in \mathbb{N}$. Let $\lambda > 0$ be such that $\mathbf{M}_n(\psi) \preceq \lambda \mathbf{M}_n(\phi)$ for all $n \in \mathbb{N}$. Then $\psi \leq \lambda \phi$ and there exists $0 \leq h \in L^\infty(\Omega, \phi)$ such that*

$$(1.1) \quad d\psi = h(\mathbf{x}) d\phi \quad \text{and} \quad \|h\|_\infty \leq \lambda.$$

Moreover,

$$(1.2) \quad \|h\|_\infty = \inf_{\lambda} \{ \lambda : \mathbf{M}_n(\psi) \preceq \lambda \mathbf{M}_n(\phi), \quad \forall n \in \mathbb{N} \}.$$

2. MAIN RESULT

Let μ be a finite Borel probability measure on Ω with density $f \in L^\infty(\Omega)$, i.e., $d\mu = f(\mathbf{x}) 1_\Omega(\mathbf{x}) d\mathbf{x}$ and $\int_\Omega f(\mathbf{x}) d\mathbf{x} = 1$.

Next, given $\boldsymbol{\xi} \in \mathbb{R}^d$ let $\mathbf{B}(\boldsymbol{\xi}; \delta) := \{\mathbf{x} \in \mathbb{R}^d : \|\mathbf{x} - \boldsymbol{\xi}\| \leq \delta\}$, and let $d\phi_\xi = \frac{1}{\text{vol}(\mathbf{B}(\boldsymbol{\xi}, \delta))} 1_{\mathbf{B}(\boldsymbol{\xi}, \delta)} d\mathbf{x}$. Given $h \in L^\infty(\Omega)$ let $\|h\|_\infty^{\boldsymbol{\xi}, \delta}$ be the infinity-norm on $\Omega \cap \mathbf{B}(\boldsymbol{\xi}, \delta)$.

Consider the extended Christoffel function:

$$(2.1) \quad \Lambda_n^\mu(\phi_\xi) := \min_{p \in \mathbb{R}^{[\mathbf{x}]_n}} \left\{ \int_\Omega p^2 d\mu : \int_{\mathbf{B}(\boldsymbol{\xi}, \delta)} p^2 d\phi_\xi = 1 \right\},$$

and observe that if $\delta = 0$ then $\Lambda_n^\mu(\phi_\xi) = \Lambda_n^\mu(\delta_\xi)$ coincides with the standard Christoffel function $\boldsymbol{\xi} \mapsto \Lambda_n^\mu(\boldsymbol{\xi})$.

Lemma 2.1. *Let $\Lambda_n^\mu(\phi_\xi)$ be as in (2.1) and $\boldsymbol{\xi} \in \text{int}(\Omega)$. Then:*

$$(2.2) \quad \Lambda_n^\mu(\phi_\xi) = \lambda_{\min}(\mathbf{M}_n(\mu), \mathbf{M}_n(\phi_\xi)), \quad \forall n \in \mathbb{N}.$$

Equivalently:

$$(2.3) \quad \begin{aligned} \Lambda_n^\mu(\phi_\xi)^{-1} &= \lambda_{\max}(\mathbf{M}_n(\phi_\xi), \mathbf{M}_n(\mu)) \\ &= \inf_{\lambda} \{ \lambda : \mathbf{M}_n(\phi_\xi) \preceq \lambda \mathbf{M}_n(\mu) \} \leq \|f\|_{\infty}^{\xi, \delta}, \end{aligned}$$

and moreover,

$$(2.4) \quad \lim_{n \rightarrow \infty} \Lambda_n^\mu(\phi_\xi)^{-1} = \|f\|_{\infty}^{\xi, \delta}.$$

Proof. Observe that (2.1) reads

$$(2.5) \quad \Lambda_n^\mu(\phi_{\xi, \delta}) = \min_{\mathbf{p} \in \mathbb{R}^{s(n)}} \{ \langle \mathbf{M}_n(\mu), \mathbf{p}\mathbf{p}^T \rangle : \langle \mathbf{M}_n(\phi_\xi), \mathbf{p}\mathbf{p}^T \rangle = 1 \}.$$

By Barvinok \square the semidefinite relaxation

$$\theta := \min_{\mathbf{X} \succeq 0} \{ \langle \mathbf{M}_n(\mu), \mathbf{X} \rangle : \langle \mathbf{M}_n(\phi_\xi), \mathbf{X} \rangle = 1 \},$$

of (2.5) has a rank-one optimal solution $\mathbf{X}^* = \mathbf{q}\mathbf{q}^T \succeq 0$ for some $\mathbf{q} \in \mathbb{R}^{s(n)}$, and therefore $\mathbf{x} \mapsto q(\mathbf{x}) := \mathbf{q}^T \mathbf{v}_n(\mathbf{x}) \in \mathbb{R}[\mathbf{x}]_n$ is an optimal solution of (2.1), which also yields $\theta = \Lambda_n^\mu(\phi_\xi)$. As $0 \prec \mathbf{X} := \mathbf{I}/\text{trace}(\mathbf{M}_n(\phi_\xi))$ is a feasible solution, Slater's condition holds and therefore by standard conic duality:

$$\Lambda_n^\mu(\phi_\xi) = \sup_{\lambda} \{ \lambda : \lambda \mathbf{M}_n(\phi_\xi) \preceq \mathbf{M}_n(\mu) \} = \lambda_{\min}(\mathbf{M}_n(\mu), \mathbf{M}_n(\phi_\xi)),$$

which yields (2.2) and also the two equalities in (2.3). To get the inequality just use Lemma 1.1.

Next by (2.3), the sequence $(\Lambda_n^\mu(\phi_\xi)^{-1})_{n \in \mathbb{N}}$ is monotone increasing and is bounded above. Therefore $\lim_{n \rightarrow \infty} \Lambda_n^\mu(\phi_\xi)^{-1} = \rho^*$ for some $\rho^* \leq \|f\|_{\infty}^{\xi, \delta}$. But then by construction, $\mathbf{M}_n(\phi_\xi) \preceq \rho^* \mathbf{M}_n(\mu)$ for all $n \in \mathbb{N}$, which implies $\rho^* \geq \|f\|_{\infty}^{\xi, \delta}$, and which in turn yields the desired result. \square

As an immediate consequence, we obtain the following result.

Lemma 2.2. *For every $n \in \mathbb{N}$ define:*

$$(2.6) \quad \Lambda_n^\mu(\phi_\xi) := \min_{p \in \mathbb{R}[\mathbf{x}]_n} \left\{ \int_{\Omega} p^2 d\mu : \int_{\mathbf{B}(\xi, 1/n)} p^2 d\phi_\xi = 1 \right\}.$$

If f is piecewise continuous and $\xi \in \text{int}(\Omega)$ is not a point of discontinuity then:

$$(2.7) \quad \liminf_{n \rightarrow \infty} \Lambda_n^\mu(\phi_\xi) \geq \frac{1}{f(\xi)}.$$

Proof. If $\xi \in \text{int}(\Omega)$ then $\mathbf{B}(\xi, 1/n) \subset \text{int}(\Omega)$ provided that n is sufficiently large, say $n \geq n_0$. So let $\Lambda_n^\mu(\phi_\xi)$ be as in (2.6) with $n \geq n_0$. If f is not a point of discontinuity then $\|f\|_{\infty}^{\xi, 1/n} \downarrow f(\xi)$ as n increases. Moreover by (2.3):

$$\limsup_{n \rightarrow \infty} \Lambda_n^\mu(\phi_\xi)^{-1} \leq \limsup_{n \rightarrow \infty} \|f\|_{\infty}^{\xi, 1/n} = f(\xi),$$

which yields the desired result (2.7). \square

In topological data analysis, they analyze clouds \mathcal{N} of finitely many points by approximating the cloud with $\bigcup_{\xi \in \mathcal{N}} \mathbf{B}(\xi, \varepsilon)$.

Notice that from KKT-optimality conditions, an optimal solution \mathbf{p} of (2.1) satisfies: $\mathbf{M}_n(\mu) \mathbf{p} = \Lambda_n^\mu(\phi_\xi) \mathbf{M}_n(\phi_\xi) \mathbf{p}$ and so

$$\mathbf{p} = \Lambda_n^\mu(\phi_\xi) \mathbf{M}_n(\mu)^{-1} \int_{\Omega} \mathbf{v}_n(\mathbf{y}) \underbrace{\mathbf{v}_n(\mathbf{y})^T \mathbf{p}}_{=p(\mathbf{y})} d\phi_\xi(\mathbf{y}).$$

and using $\mathbf{v}_n(\mathbf{x})^T \mathbf{M}_n(\mu)^{-1} \mathbf{v}_n(\mathbf{y}) = \sum_{\alpha} P_\alpha(\mathbf{x}) P_\alpha(\mathbf{y})$, yields:

$$\begin{aligned} \mathbf{x} \mapsto p(\mathbf{x}) = \mathbf{v}_n(\mathbf{x})^T \mathbf{p} &= \Lambda_n^\mu(\phi_\xi) \sum_{\alpha \in \mathbb{N}_n^d} P_\alpha(\mathbf{x}) \int_{\mathbf{B}(\xi, \delta)} P_\alpha(\mathbf{y}) p(\mathbf{y}) d\phi_\xi(\mathbf{y}). \\ &= \Lambda_n^\mu(\phi_\xi) \int_{\mathbf{B}(\xi, \delta)} \sum_{\alpha \in \mathbb{N}_n^d} P_\alpha(\mathbf{x}) P_\alpha(\mathbf{y}) p(\mathbf{y}) d\phi_\xi(\mathbf{y}) \\ &= \Lambda_n^\mu(\phi_\xi) \int_{\mathbf{B}(\xi, \delta)} K(\mathbf{x}, \mathbf{y}) p(\mathbf{y}) d\phi_\xi(\mathbf{y}) \end{aligned}$$

So the optimal solution $p \in \mathbb{R}[\mathbf{x}]_n$ is reproduced by the kernel $K(\mathbf{x}, \mathbf{y})$ on $\mathbf{B}(\xi, \delta)$ (whereas K is a RKHS on $L^2(\mu)$ (and not on $L^2(\phi_\xi)$). Moreover:

$$\Lambda_n^\mu(\phi_\xi)^{-1} = \sum_{\alpha \in \mathbb{N}_n^d} \left(\int_{\mathbf{B}(\xi, \delta)} P_\alpha(\mathbf{y}) p(\mathbf{y}) d\phi_\xi(\mathbf{y}) \right)^2,$$

which coincides with $\Lambda_n^\mu(\xi)^{-1} = K(\xi, \xi)$ when $\phi_\xi = \delta_\xi$ (as then $p(\xi) = 1$).

REFERENCES

- [1] F. CHAZAL, D. COHEN-STEINER, Q. MÉRIGOT Geometric Inference for Probability Measures, *FoCM* **11** (2011), pp. 733–751.